


I hereby certify that this correspondence is enclosed in an envelope being deposited with the United States Postal Service as first class mail under 37 CFR 1.10 on the date indicated above and is addressed to:

PATENT
Attorney Docket No.: 015280-290100US
Client Ref. No.: E-171-96/0

Assistant Commissioner for Patents
Art Unit 1644
Washington, D.C. 20231.



TOWNSEND AND TOWNSEND AND CREW LLP

On: August 27, 2001

By: H. Munler
H. Munler

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

KLIMPEL *et al.*

Application No.: 08/937,276

Filed: September 15, 1997

For: TARGETING ANTIGENS TO THE
MHC CLASS I PROCESSING PATHWAY
WITH AN ANTHRAX TOXIN FUSION
PROTEIN

Examiner: Ronald B. Schwadron, Ph.D.

Art Unit: 1644

DECLARATION OF DR. JAY A.
BERZOFKY UNDER 37 C.F.R. § 1.132

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

I, Jay A. Berzofsky, M.D., Ph.D., being duly warned that willful false statements and the like are punishable by fine or imprisonment or both (18 U.S.C. § 1001), and may jeopardize the validity of the patent application or any patent issuing thereon, state and declare as follows:

1. All statements herein made of my own knowledge are true, and statements made on information or belief are believed to be true and correct.

2. I am the Chief of the Molecular Immunogenetics and Vaccine Research Section, Metabolism Branch, Center for Cancer Research, National Cancer Institute, National Institutes of Health. I have been in this position since 1987. I am currently the Chair for the Vaccine Working Group for the National Cancer Institute.

3. I received a bachelor's degree in chemistry, summa cum laude, from Harvard University in 1967. I received a Ph.D. and an M.D. degree from the Albert Einstein School of Medicine in 1972 and 1973, respectively. I trained in medicine at Massachusetts General Hospital, and as a postdoctoral fellow in the laboratory of Nobel prize winner Dr. Christian B. Anfinsen at the National Institutes of Health. Prior to 1987, I was a Senior Investigator at the National Institutes of Health. I am also a past president of the American Society for Clinical Investigation. I am an author of 330 published or in press scientific papers and books. In addition, I am an author of chapters in several major immunology textbooks. A copy of my curriculum vitae and bibliography is attached hereto as Exhibit A.

4. The present invention relates to the surprising discovery that the anthrax toxin system can be used to deliver full length proteins to the cell cytosol for processing and presentation by MHC class I molecules to cytotoxic T cells (CTLs). In particular, the invention provides the first evidence that an anthrax toxin fusion protein comprising a full length protein-anthrax lethal factor (LF) fusion bound to anthrax protective antigen (PA) is translocated into the cell, and that the full length protein is efficiently processed into multiple epitopes by the MHC class I processing pathway and presented by MHC class I molecules to CTLs. The use of full length protein has the advantage of providing multiple epitopes that are recognized by more than one MHC class I allele. Moreover, the anthrax toxin fusion proteins of the present invention, comprising the full length protein, are more potent for sensitizing CTLs than corresponding peptide epitopes.

5. I am a named inventor on the above-referenced patent application. I have read and am familiar with the contents of this patent application. In addition, I have read the Office Action, dated February 27, 2001, received in the present case. It is my

understanding that the Examiner is concerned that the claims are obvious over Milne *et al.*, Arora *et al.*, or Leppla *et al.* in view of Donnolly *et al.* and EP 0 532 090. In particular, the Examiner asserts that it would have been obvious to one of ordinary skill in the art at the time of the invention to use an anthrax toxin fusion protein to deliver a protein to a cell, as taught by Milne *et al.*, Leppla *et al.*, or Arora *et al.*, for processing of the protein via the cytosolic MHC class I pathway and induction of a CTL response to the protein, as taught in Donnelly *et al.*

6. This declaration is provided to demonstrate that, at the time of the invention, one of skill in the art would not have had a reasonable expectation of success using the anthrax toxin fusion proteins of Milne *et al.*, Arora *et al.*, or Leppla *et al.* in combination with the methods of inducing an immune response described by Donnolly *et al.* and EP 0 532 090. Furthermore, one of skill in the art would not have been motivated to substitute an anthrax toxin fusion protein for the *Pseudomonas* toxin (PE) fusion protein used in the methods of Donnolly *et al.* and EP 0 532 090 to practice the invention. Moreover, this declaration is provided to demonstrate that, using the anthrax toxin fusion proteins of the present invention to induce an immune response, the inventors achieved results that are more effective than and superior to those of the closest prior art.

7. In my scientific opinion, Milne *et al.*, Arora *et al.*, and Leppla *et al.* teach methods for cellular delivery of small fragments of proteins or peptide epitopes, rather than full length proteins, using anthrax toxins comprising PA and an LF fusion protein. Anthrax lethal toxin is a binary bacterial toxin comprising two proteins, LF and PA. PA binds to a cellular receptor and is cleaved, revealing an LF binding site. LF binds to PA, forming anthrax toxin, which is translocated into the cell. However, as described in detail below, these references fail to teach the use of anthrax toxin fusion protein to elicit a T cell response, by delivering proteins or peptides to a cytosolic MHC class I pathway for epitope processing and cell surface presentation.

For example, Milne *et al.* teaches compositions where the catalytic domain of diphtheria toxin A chain is fused to LF and translocated into a cell, via PA binding. Arora *et al.* teaches compositions where the catalytic domain of *Pseudomonas* exotoxin A is fused to LF and translocated into a cell, via PA binding. Leppla *et al.* teaches generic compositions where peptides or protein fragments are fused to LF for cellular translocation, via PA binding.

Each of these references therefore discloses methods of internalizing protein fragments or peptide epitopes, for delivery of an activity to a cell, e.g., a cytotoxic activity. None of these references teach that anthrax toxin can be used to deliver a protein to a cytosolic MHC class I processing pathway, for induction of an immune response.

Furthermore, the teachings of Milne *et al.*, Arora *et al.*, and Leppla *et al.* at most only propose the use of anthrax toxins to deliver full length proteins to cells. Also, Arora *et al.* at most only proposes that anthrax toxin could be used to present peptides to MHC class I molecules. However, as described below, the art teaches away from the use of bacterial toxins to present peptides to MHC class I molecules, let alone full length proteins. These teachings therefore do not provide any indication as to what parameters are critical, nor provide any direction as to which of the many possible choices for making and using anthrax toxins for delivery of full length proteins to the cytosol for processing and presentation by MHC class I molecules to CTLs. None of these references would provide one of skill in the art with a reasonable expectation of success for practicing the claimed invention.

8. In my scientific opinion, Donnelly *et al.* and EP 0 532 090 teach methods of inducing an immune response using *Pseudomonas* exotoxin ("PE") fused to a peptide epitope (*i.e.*, peptide epitopes from influenza A matrix protein or nucleoprotein). PE is a single subunit bacterial toxin, rather than a binary toxin like anthrax. These references therefore teach neither the use of binary toxins nor the use of full length proteins to induce an immune response. The references merely describe attempts to use such PE fusion proteins to translocate peptide epitopes into a cell for presentation by the MHC class I pathway.

Moreover, the same authors in a separate publication by Ulmer *et al.* (further discussed below) demonstrate that PE toxin fusion proteins are likely processed via an alternative, endosomal processing pathway that is less efficient than the MHC class I cytosolic pathway. Therefore, one skilled in the art would conclude that such bacterial toxin fusion proteins do not deliver antigen to the MHC class I cytosolic pathway. This reference thus teaches away from using bacterial toxin fusion proteins for processing of an antigen and presentation of an antigen by MHC class I molecules.

In addition, these references teach the administration of a 12-mer amino acid epitope fused to PE toxin. Due to its size, such an epitope does not require cellular processing in order to be displayed by MHC class I molecules. Prior to the time of the present invention, the disadvantages associated with the translocation of large proteins (*e.g.*, inefficient translocation) and the advantages associated with the use of peptide epitopes (*e.g.*, efficient translocation and generation of T cells directed to a single epitope) were known to one of skill

in the art. Consistent with this understanding, Donnelly *et al.* and EP 0 532 090 teach the use of bacterial toxin fusion proteins encoding peptide epitopes, and not full length proteins. These references therefore fail to provide the requisite motivation to one of skill in the art to use anthrax toxin fusion proteins for delivery of a full length protein to a cell for processing and presentation by MHC class I.

9. The present application provides the first evidence that a bacterial toxin system (anthrax toxin) can be used to introduce a full length protein into the cytosol for processing via the MHC class I pathway and presentation by MHC class I molecules to CTLs. This processing of the full length protein and presentation to CTLs, via the MHC class I pathway, was demonstrated by treating antigen-presenting cells with lactacystin, which inhibits proteasome function required for cytosolic MHC class I processing (*see* specification, page 3, lines 7-11). In addition, the evidence demonstrates that the anthrax toxins of the present invention, fused to full length proteins, are more potent for sensitizing CTLs than a peptide epitope from the same full length protein (*see* Goltez *et al.*, *PNAS USA* 94:12059-12064 (1997), page 12061, previously submitted on May 24, 1999 as Appendix A).

10. The present inventors are therefore the first to show that a full length protein, fused to LF and translocated into a cell by anthrax toxin, is processed by the cytosolic MHC class I pathway and presented by MHC class I molecules to CTLs. In addition, the fusion proteins of the present invention are more potent for sensitizing CTLs than peptide epitopes and thus provide superior results over that taught in the prior art. In contrast, the prior art describes only the use of bacterial toxins to translocate protein fragments or epitopes into cells, for delivery of, e.g., a cytotoxic activity. In addition, the prior art teaches that bacterial toxin fusion proteins are likely processed by an alternative endocytic processing pathway, and not the cytosolic MHC class I processing pathway that results in the efficient presentation of a processed protein by MHC class I molecules to CTLs. Thus, the prior art teaches away from the claimed invention. But, even if one of skill in the art was motivated to try this approach, undue experimentation would be required to practice the claimed invention in view of the teachings of the prior art. Therefore, it is my scientific opinion that these prior studies would have provided neither motivation nor a reasonable expectation of success to use anthrax toxin to introduce a full length protein into a cell and elicit a CTL immune response, by processing the full length protein via the cytosolic MHC class I pathway, and presenting the resulting

epitopes by MHC class I molecules. Moreover, in view of the prior art teachings, our results are unexpected and surprisingly effective.

11. In the Office Action, the Examiner asserts that “the ability of the protein-LF PA binary complex to translocate the protein into the cytosolic MHC class I processing pathway is an inherent property of the complex.” I disagree with the Examiner’s assertion for the following reasons. Prior to the time of the invention, such properties of anthrax toxin proteins would have been unpredictable and unexpected. At that time, the prior art taught that peptide epitopes delivered to a cell using bacterial fusion toxin proteins likely bind to preexisting MHC class I molecules that are recycled and internalized via an endocytic pathway, rather than being processed by the more efficient cytosolic MHC class I pathway. The prior art provides no information regarding processing of bacterial toxin fusion proteins via the cytosolic MHC class I pathway. Moreover, the prior art taught that large proteins were inefficiently translocated into the cell, if at all. The present inventors are the first to demonstrate that anthrax toxins can be used to deliver whole proteins to the cell cytosol for processing and presentation by MHC class I molecules to CTLs. Prior to this discovery, such properties and uses of bacterial toxin proteins for delivery of whole proteins to cells would have been unpredictable and unexpected.

12. The Examiner further asserts that Arora *et al.* teach that anthrax toxin fusion proteins are translocated into the cell and induce a CTL response to the heterologous protein fused to LF; and that the teachings of Ulmer *et al.* support the Examiner’s assertions. For the following reasons, I disagree with the Examiner’s assertion.

As mentioned above, the teachings of Arora *et al.* are directed to cellular delivery of small fragments of proteins or peptide epitopes and at most only propose the use of whole proteins. Arora *et al.* do not provide any indication as to what parameters are critical, nor provide any direction as to which of the many possible choices for making and using anthrax toxin fusion proteins for delivery of whole proteins to the cytosol for processing via the MHC class I pathway and presentation by MHC class I molecules to CTLs.

Moreover, the teachings of Ulmer *et al.* relied on by the Examiner actually - teach away from the claimed invention. The conclusion of the Ulmer *et al.* study is that the PE


fusion protein is not translocated into the cytosolic MHC class I pathway, but rather enters an endosomal pathway. This reference therefore teaches away from the use of bacterial toxin fusion proteins for introducing whole proteins into the cytosolic MHC class I pathway for subsequent processing and presentation by MHC class I molecules.

Further, the Examiner points out that such experiments by Ulmer *et al.* "do not exclude that under physiological conditions . . . translocation into the cytosolic MHC class I processing does not occur" (emphasis added). Merely teaching that the experiments do not exclude what is not known fails to provide support or a rational basis for concluding that the prior art teaches or suggests the translocation of bacterial toxin fusion proteins into the cytosolic MHC class I processing pathway.

Therefore, the teachings of Ulmer *et al.* do not support the Examiner's assertion that Arora *et al.* teaches the use of anthrax toxin fusion proteins for delivery of whole proteins to the cytosol for processing by the MHC class I processing pathway and presentation by MHC class I molecules to CTLs.

13. In conclusion, the present application provides the first evidence that full length proteins fused to anthrax toxin are translocated into the cell, enter the cytosolic MHC class I processing pathway, and are subsequently presented by MHC class I molecules to CTLs. This evidence is contrary to the teachings of the prior art, which suggest that bacterial toxin fusion proteins are likely processed via an endocytic pathway. Moreover, the anthrax toxin fusion proteins of the present invention, comprising a full length protein, are more potent for sensitizing CTLs than peptide epitopes. Thus, using the anthrax toxin fusion proteins of the present invention, the inventors achieved results that are more effective than and superior to those of the closest prior art. In view of the prior art, these results are unobvious and unexpected, because the prior art teaches the advantages and use of small fragments of proteins and peptide epitopes over full length proteins, and teaches disadvantages associated with the translocation of full length proteins. Thus, it is my scientific opinion that the teachings of Milne *et al.*, Arora *et al.*, or Leppla *et al.* in view of Donnolly *et al.* and EP 0 532 090 do not provide one of skill in the art with a reasonable expectation of success, nor motivate one of skill in the art to practice the invention.

- Date: 24 August, 2001

By: 

CURRICULUM VITAE

Name: Jay Arthur Berzofsky

Date and Place of Birth: April 13, 1946, Baltimore, Maryland

Marital Status: Married to Sharon M. Miller, June 27, 1969; two children
Alexander, April 30, 1974, and Marcus, May 27, 1976

Education:

1967 - A.B., Harvard University (Summa Cum Laude in Chemistry)
1971 - Ph.D., Albert Einstein College of Medicine, Molecular Biology
1973 - M.D., Albert Einstein College of Medicine, Medical Scientist
Training Program

Brief Chronology of Employment:

1973 - 1974	Medical Internship (Straight Medicine), Massachusetts General Hospital, Boston, Massachusetts
1974 - 1976	Research Associateship, Laboratory of Chemical Biology National Institute of Arthritis, Metabolism, and Digestive Diseases, National Institutes of Health
1976 - 1979	Investigator ("Expert"), Metabolism Branch, National Cancer Institute, National Institutes of Health
1979 - 1987	Senior Investigator, Metabolism Branch, National Cancer Institute, National Institutes of Health
1987 - Date	Chief, Molecular Immunogenetics and Vaccine Research Section, Metabolism Branch, National Cancer Institute, National Institutes of Health

Honors/Awards:

Detur Prize, Harvard University, 1964
Harvard College Scholarship, Harvard University, 1964
Phi Beta Kappa, Junior Year, Harvard University, 1966
Summa Cum Laude in Chemistry, Harvard University, 1967
Sophia Freund Prize for Graduate with Highest Cumulative Grade Point
Average, Harvard College, 1967
NIH Special Achievement Award, 1982
Hollister - Stier's Distinguished Lectureship, Washington State
University, 1986
J. W. McLaughlin Fund Distinguished Contributions to Immunology Lectureship,
University of Texas Medical School, Galveston, 1987

U. S. Public Health Service Superior Service Award, 1988
 31st Michael Heidelberger Award and Lecture, Columbia University, 1992
 McLaughlin Visiting Professorship, University of Texas Medical School,
 Galveston, 1992
 American Society for Clinical Investigation, President 1993-94
 Fellow of the American Association for the Advancement of Science, 1995
 Cytokine Interest Group Best Paper of 2000 Award to fellow in lab, 2001

Professional Society Memberships:

Association of Harvard Chemists, 1967 - present
 New York Academy of Sciences, 1971 - present
 American Association of Immunologists, 1977 - present
 Undersea Medical Society, 1978 - 1988
 American Federation for Clinical Research, 1979 - present
 American Society of Biological Chemists, 1980 - present
 American Society for Clinical Investigation, 1983 - present,
 Secretary-Treasurer, 1989 - 1992
 President-elect, 1992-1993
 President, 1993-94
 Association of American Physicians, 1990 - present

Editorial Positions:

Associate Editor, *Journal of Immunology*, 1980 - 1984
 Editorial Advisory Board, *Journal of Molecular and Cellular Immunology*, 1983 -
 present
 Advisory Editor, *Molecular Immunology*, 1985 - 1988
 Editorial Board, *Peptide Research*, 1987 - present
 Transmitting Editor, *International Immunology*, 1988 - present
 Editorial Board, *Journal of Human Virology*, 1997-present
 Consulting Editor, *Journal of Clinical Investigation*, 1998-present
 Section Editor, *Journal of Clinical Immunology*, 2002-present

Professional Committees:

American Association of Immunologists, Membership Committee, 1981 - 1988
 American Association of Immunologists, Chairman of Membership Committee,
 1983 - 1988
 NIH Clinical Center Compensable Events Committee, 1982 - present
 American Society for Clinical Investigation, Council, 1989-1994
 NCI Division of Clinical Sciences Promotion and Tenure Committee, 1995-
 present.
 NCI Division of Clinical Sciences Research Advisory Group, 1995-present
 NCI Director's Intramural Advisory Board, 1997-99
 NIH AIDS Vaccine Research Center Steering Committee, 1997-present
 NIH Search Committee for Director of Office of AIDS Research, 1997-98
 NIAID Malaria Vaccine Task Force, 1998-present
 NCI Vaccine Working Group, Chairman/Organizer, 1998-present
 NCI/CCR Immunology Faculty Steering Committee, 2001-present
 NCI/CCR HIV & Virology Faculty Steering Committee, 2001-present

Military Service:

Commissioned Corps, United States Public Health Service, 1974 - 1976

Other Research Experience:

Summers, 1962 - 1965 Research Assistant, Pediatric Research Unit (H. M. Nitowsky), Sinai Hospital, Baltimore, Maryland
Summer, 1966 Research Assistant, Organic Synthesis Laboratory (C. H. Robinson), Department of Pharmacology, Johns Hopkins School of Medicine, Baltimore, Maryland
Summer, 1967 Visiting Scientist, Laboratoire d'Enzymologie (G. N. Cohen), Centre National de la Recherche Scientifique, Gif-sur-Yvette, France

Medical Licensure: Maryland and Massachusetts

Major Outside Activities

Medimmune, Inc.—Scientific Founder and Chair, Scientific Advisory Board, 1989-present
 Magainin Pharmaceuticals, Inc.—Member, Scientific Advisory Board, 1991-97
 Diacrin, Inc.—Member, Scientific Advisory Board, 1993-present
 Pharmadyne, Inc.—Scientific Co-Founder and Chair, Scientific Advisory Board, 1997-present
 Boston University Community Technology Fund—Consultant, 1997-present
 Health Care Ventures, Inc.—consultant, 1998-present
 EMD Pharmaceuticals, Inc.—consultant, 2000-present
 Epivax, Inc.—Member, Scientific Advisory Board, 2000-present

Major areas of research:

1. Molecular basis of antigen recognition by T lymphocytes
2. Processing of antigens and their presentation by major histocompatibility molecules
3. Structure of antigenic sites on protein antigens
4. Genetic regulation of the immune response
5. Design and development of artificial vaccines based on immunological principles and peptide synthesis or recombinant DNA technology
6. AIDS vaccines and diagnostic techniques
7. Malaria vaccines
8. Cancer vaccines
9. Antigen-antibody interactions
10. Structure-function relationships in proteins and protein conformation.
11. Regulation of tumor immunosurveillance and T cell function by cytokines
12. Mucosal immunity and vaccines

BIBLIOGRAPHY

Jay Arthur Berzofsky

1. Nitowsky, H.M., L. Matz, and J.A. Berzofsky. 1966. Studies on oxidative drug metabolism in the full-term newborn infant. *J. Pediatrics* 69:1139-1149.
2. Berzofsky, J.A., J. Peisach, and W.E. Blumberg. 1971. Sulfheme proteins. I. Optical and magnetic properties of sulfmyoglobin and its derivatives. *J. Biol. Chem.* 246:3367-3377.
3. Peisach, J., J.A. Berzofsky, and W.E. Blumberg. 1973. Electronic control of oxygen binding to heme proteins. In *Proceedings of the Second International Symposium on Oxidases and Related Oxidation-Reduction Systems*. T.E. King, H.S. Mason, and M. Morrison, editors. University Park Press, Baltimore. 265-278.
4. Berzofsky, J.A. 1971. The nature of sulfmyoglobin: Chemical, physical, and oxygen-binding properties. Albert Einstein College of Medicine, New York.
5. Berzofsky, J.A., J. Peisach, and W.E. Blumberg. 1971. Sulfheme proteins II. The reversible oxygenation of ferrous sulfmyoglobin. *J. Biol. Chem.* 246:7366-7372.
6. Berzofsky, J.A., J. Peisach, and J.O. Alben. 1972. Sulfheme proteins. III. Carboxysulfmyoglobin: The relation between electron withdrawal from iron and ligand binding. *J. Biol. Chem.* 247:3774-3782.
7. Berzofsky, J.A., J. Peisach, and B.L. Horecker. 1972. Sulfheme proteins. IV. The stoichiometry of sulfur incorporation and the isolation of sulfhemin, the prosthetic group of sulfmyoglobin. *J. Biol. Chem.* 247:3783-3791.
8. Berzofsky, J.A., A.N. Schechter, and H. Kon. 1976. Does Freund's adjuvant denature protein antigens? EPR studies of emulsified hemoglobin. *J. Immunol.* 116:270-273.
9. Berzofsky, J.A., J.G. Curd, and A.N. Schechter. 1976. Probability analysis of the interaction of antibodies with multideterminant antigens in radioimmunoassay: application to the amino terminus of the beta chain of hemoglobin S. *Biochem* 15:2113-2121.
10. Berzofsky, J.A., A.N. Schechter, G.M. Shearer, and D.H. Sachs. 1977. Genetic control of the immune response to staphylococcal nuclease III. Time course and correlation between the response to native nuclease and the response to its polypeptide fragments. *J. Exp. Med.* 145:111-112.
11. Berzofsky, J.A., A.N. Schechter, G.M. Shearer, and D.H. Sachs. 1977. Genetic control of the immune response to staphylococcal nuclease IV. H-2-linked control of the relative proportions of antibodies produced to different determinants of native nuclease. *J. Exp. Med.* 145:123-145.

12. Sachs, D.H., J.A. Berzofsky, C.G. Fathman, D.S. Pisetsky, A.N. Schechter, and R.H. Schwartz. 1976. The immune response to staphylococcal nuclease: A probe of cellular and humoral antigen specific receptors. *Cold Spring Harbor Symp. Quant. Biol.* 41:295-306.
13. Berzofsky, J.A., R.H. Schwartz, A.N. Schechter, and D.H. Sachs. 1978. H-2 linked control of the antibody and cellular immune response to nuclease at the level of individual regions of the molecule. In *Ir Genes and Ia Antigens* (Proceedings of the Third Ir Gene Workshop, December, 1976, Asilomar, CA). H.O. McDevitt, editor. Academic Press, New York. 423-431.
14. Berzofsky, J.A. 1978. Genetic control of the immune response to mammalian myoglobins in mice I. More than one I-region gene in H-2 controls the antibody response. *J. Immunol.* 120(2):360-369.
15. Pisetsky, D.S., J.A. Berzofsky, and D.H. Sachs. 1978. Genetic control of the immune response to staphylococcal nuclease. VII. Role of non-H-2-linked genes in the control of the anti-nuclease antibody response. *J. Exp. Med.* 147(2):396-408.
16. Berzofsky, J.A. 1978. The assessment of antibody affinity from radioimmunoassay. *Clinical Chem.* 24(3):419-421.
17. Berzofsky, J.A. 1978. Genetic control of the antibody response to sperm whale myoglobin in mice. *Adv. Exp. Med. Biol.* 98:225-240.
18. Berzofsky, J.A., D.S. Pisetsky, R.H. Schwartz, A.N. Schechter, and D.H. Sachs. 1978. Genetic control of the immune response to staphylococcal nuclease in mice. *Adv. Exp. Med. Biol.* 98:241-258.
19. Schwartz, R.H., J.A. Berzofsky, C.L. Horton, A.N. Schechter, and D.H. Sachs. 1978. Genetic control of the T-lymphocyte proliferative response to staphylococcal nuclease. Evidence for multiple MHC-linked Ir gene control. *J. Immunol.* 120:1741-1749.
20. Sachs, D.H., J.A. Berzofsky, D.S. Pisetsky, and R.H. Schwartz. 1978. Genetic control of the immune response to staphylococcal nuclease. In *Seminars in Immunopathology* Vol.I. B. Benacerraf, editor. Springer-Verlag, New York. 51-83.
21. Berzofsky, J.A., L.K. Richman, and D.J. Killion. 1979. Distinct H-2-linked Ir genes control both antibody and T cell responses to different determinants on the same antigen, myoglobin. *Proc. Natl. Acad. Sci. U. S. A.* 76:4046-4050.
22. Richman, L.K., R.J. Klingenstein, J.A. Richman, W. Strober, and J.A. Berzofsky. 1979. The murine Kupffer cell. I. Characterization of the cell serving accessory function in antigen-specific T cell proliferation. *J. Immunol.* 123:2602-2609.

23. Richman, L.K., W. Strober, and J.A. Berzofsky. 1980. Genetic control of the immune response to myoglobin: III. Determinant-specific, two Ir gene phenotype is regulated by the genotype of reconstituting Kupffer cells. *J. Immunol.* 124:619-624.
24. Berzofsky, J.A. 1980. Immune response genes in the regulation of mammalian immunity. In *Biological Regulation and Development*. R.F. Goldberger, editor. Plenum Press, New York. 467-594.
25. Berzofsky, J.A., G. Hicks, J. Fedorko, and J. Minna. 1980. Properties of monoclonal antibodies specific for determinants of a protein antigen, myoglobin. *J. Biol. Chem.* 255:11188-11191.
26. Berzofsky, J.A., L.K. Richman, and W. Strober. 1982. Determinant-specific antigen presentation by liver Kupffer cells under control of H-2-linked Ir genes. In *Recent Advances in Mucosal Immunity*. W. Strober, L.A. Hanson, and K.W. Sell, editors. Raven Press, New York. 213-223.
27. Berzofsky, J.A., and A.N. Schechter. 1981. The concepts of crossreactivity and specificity in immunology. *Molec. Immunol.* 18:751-763.
28. Berzofsky, J.A., and L.K. Richman. 1981. Genetic control of the immune response to myoglobins. IV. Inhibition of determinant-specific Ir-gene-controlled antigen presentation and induction of suppression by pretreatment of presenting cells with anti-Ia antibodies. *J. Immunol.* 126:1898-1904.
29. Berzofsky, J.A., G.K. Buckenmeyer, G. Hicks, F.R.N. Gurd, R.J. Feldmann, and J. Minna. 1982. Topographic antigenic determinants recognized by monoclonal antibodies to sperm whale myoglobin. *J. Biol. Chem.* 257:3189-3198.
30. Berzofsky, J.A., D.S. Pisetsky, D.J. Killion, G. Hicks, and D.H. Sachs. 1981. Ir genes of different high responder haplotypes for staphylococcal nuclease are not allelic. *J. Immunol.* 127:2453-2455.
31. Berzofsky, J.A., G.K. Buckenmeyer, G. Hicks, D.J. Killion, I. Berkower, Y. Kohno, M.A. Flanagan, M.R. Busch, R.J. Feldmann, J. Minna, and F.R.N. Gurd. 1983. Topographic antigenic determinants detected by monoclonal antibodies to myoglobin. In *Protein Conformation as Immunological Signal*. F. Celada, E. Sercarz, and V. Shumaker, editors. Plenum Press, New York. 165-180.
32. Berkower, I., G.K. Buckenmeyer, F.R.N. Gurd, and J.A. Berzofsky. 1983. A possible immunodominant domain of myoglobin for murine T lymphocytes. In *Protein Conformation as Immunological Signal*. F. Celada, E. Sercarz, and V. Shumaker, editors. Plenum Press, New York. 289-302.
33. Berzofsky, J.A., G.K. Buckenmeyer, and G. Hicks. 1982. Genetics control of the immune response to myoglobins. VI. Distinct Ir genes for different myoglobins: Complementing genes in I-A and H-2D for equine myoglobin. *J. Immunol.* 128:737-741.

34. Kohno, Y., I. Berkower, J. Minna, and J.A. Berzofsky. 1982. Idiotypes of anti-myoglobin antibodies: Shared idiotypes among monoclonal antibodies to distinct determinants of sperm whale myoglobin. *J. Immunol.* 128:1742-1748.
35. Berzofsky, J.A., and I. Berkower. 1983. Antigen-Antibody Interaction. In *Fundamental Immunology*. W.E. Paul, editor. Raven Press, New York. 595-644.
36. Kohno, Y., and J.A. Berzofsky. 1982. Genetic control of the immune response to myoglobin. V. Antibody production in vitro is macrophage and T cell-dependent and is under control of two determinant-specific Ir genes. *J. Immunol.* 128:2458-2464.
37. Lando, G., J.A. Berzofsky, and M. Reichlin. 1982. Antigenic structure of sperm whale myoglobin: I. Partition of specificities between antibodies reactive with peptides and native protein. *J. Immunol.* 129:206-211.
38. Berkower, I., G.K. Buckenmeyer, F.R.N. Gurd, and J.A. Berzofsky. 1982. A possible immunodominant epitope recognized by murine T lymphocytes immune to different myoglobins. *Proc. Natl. Acad. Sci. U. S. A.* 79:4723-4727.
39. Kohno, Y., and J.A. Berzofsky. 1982. Genetic control of the immune response to myoglobins. Both low and high responder T cells tolerant to the other MHC help high but not low responder B cells. *J. Exp. Med.* 156:791-809.
40. Simmerman, H.K.B., C.C. Wang, E.M. Horwitz, J.A. Berzofsky, and F.R.N. Gurd. 1982. Semisynthesis of sperm whale myoglobin by fragment condensation. *Proc. Natl. Acad. Sci. U. S. A.* 79:7739-7743.
41. Kohno, Y., and J.A. Berzofsky. 1982. Genetic control of immune response to myoglobin. Ir gene function in genetic restriction between T and B lymphocytes. *J. Exp. Med.* 156:1486-1501.
42. Berzofsky, J.A., Y. Kohno, and H. Kawamura. 1983. Both low and high responder myoglobin-specific T cells, tolerant to F1, help high but not low responder B cells. In *Ir Genes: Past, Present, and Future*. C.W. Pierce, S.E. Cullen, J.A. Kapp, B.D. Schwartz, and D.C. Shreffler, editors. Humana Press, New Jersey. 269-275.
43. Berzofsky, J.A. 1983. T-B reciprocity: An Ia-restricted epitope-specific circuit regulating T cell-B cell interaction and antibody specificity. *Survey of Immunol. Res.* 2:223-229.
44. Berzofsky, J.A. 1984. Monoclonal antibodies as probes of antigenic structure. In *Monoclonal Antibodies as Probes of Antigenic Structure, Receptor Biochemistry and Methodology* Vol. 4. J.C. Venter, C.M. Fraser, and J.M. Lindstrom, editors. Alan R. Liss, Inc., New York. 1-19.
45. Berzofsky, J.A. 1983. T cell activation by antigen. *Immunol. Today* 4:299-301.

46. Venter, J.C., J.A. Berzofsky, J. Lindstrom, S. Jacobs, C.M. Fraser, L.D. Kohn, W. Schnieder, G.L. Greene, A.D. Strosberg, and B.F. Erlanger. 1984. Monoclonal and anti-idiotypic antibodies as probes for receptor structure and function. *Fed. Proc.* 43:2534-2539.
47. Berzofsky, J.A. 1984. Mechanisms of antigen-specific, genetically restricted, T cell-B cell interaction. *Survey of Immunol. Res.* 3:103-106.
48. Benjamin, D.C., J.A. Berzofsky, I.J. East, F.R.N. Gurd, C. Hannum, S.J. Leach, E. Margoliash, J.G. Michael, A. Miller, E.M. Prager, M. Reichlin, E.E. Sercarz, S.J. Smith-Gill, P.E. Todd, and A.C. Wilson. 1984. The antigenic structure of proteins: A reappraisal. *Annu. Rev. Immunol.* 2:67-101.
49. Berkower, I., L.A. Matis, G.K. Buckenmeyer, F.R.N. Gurd, D.L. Longo, and J.A. Berzofsky. 1984. Identification of distinct predominant epitopes recognized by myoglobin-specific T cells under control of different Ir genes and characterization of representative T-cell clones. *J. Immunol.* 132:1370-1378.
50. Kawamura, H., Y. Kohno, M. Busch, F.R.N. Gurd, and J.A. Berzofsky. 1984. A major anti-myoglobin idotype: Influence of H-2-linked Ir genes on idotype expression. *J. Exp. Med.* 160:659-678.
51. Streicher, H.Z., I.J. Berkower, M. Busch, F.R.N. Gurd, and J.A. Berzofsky. 1984. The role of antigen conformation in determining requirements for antigen processing for T cell activation. In *Regulation of the Immune System*. E. Sercarz, H. Cantor, and L. Chess, editors. Alan R. Liss, Inc., New York. 163-180.
52. Berzofsky, J.A., and H.M. Grey. 1984. MHC-restricted specific antigen presentation to different T cell sets (Th, Ts, Tc). In *Regulation of the Immune System*. E. Sercarz, H. Cantor, and L. Chess, editors. Alan R. Liss, Inc., New York. 189-193.
53. Streicher, H.Z., I.J. Berkower, M. Busch, F.R.N. Gurd, and J.A. Berzofsky. 1984. Antigen conformation determines processing requirements for T-cell activation. *Proc. Natl. Acad. Sci. U. S. A.* 81:6831-6835.
54. Berzofsky, J.A., and K. Yokomuro. 1985. Antigen presenting cells: Workshop Summary. In *Immune Regulation*. M. Feldmann, and N.A. Mitchison, editors. Humana Press, NJ. 369-374.
55. Kohno, Y., H. Kawamura, and J.A. Berzofsky. 1985. A unidirectional carrier effect. *Cellular Immunol.* 92:226-234.
56. Berzofsky, J.A. 1985. The nature and role of antigen processing in T cell activation. In *The Year in Immunology 1984-1985*. J.M. Cruse, and R.E. Lewis, Jr., editors. Karger, Basel. 18-24.

57. Morrissey, P.J., S.O. Sharrow, Y. Kohno, J.A. Berzofsky, and A. Singer. 1985. Correlation of intrathymic tolerance with intrathymic chimerism in neonatally tolerized mice. *Transplant.* 40:68-72.
58. Kawamura, H., S.A. Rosenberg, and J.A. Berzofsky. 1985. Immunization with antigen and interleukin-2 in vivo overcomes Ir genetic low responsiveness. *J. Exp. Med.* 162:381-386.
59. Berkower, I., H. Kawamura, L.A. Matis, and J.A. Berzofsky. 1985. T cell clones to two major T cell epitopes of myoglobin: Effect of I-A/I-E restriction on epitope dominance. *J. Immunol.* 135:2628-2634.
60. Delisi, C., and J.A. Berzofsky. 1985. T cell antigenic sites tend to be amphipathic structures. *Proc. Natl. Acad. Sci. U. S. A.* 82:7048-7052.
61. Berzofsky, J.A. 1985. Intrinsic and extrinsic factors in protein antigenic structure. *Science* 229:932-940.
62. Darst, S.A., C.R. Robertson, and J.A. Berzofsky. 1986. Myoglobin adsorption onto crosslinked polydimethylsiloxane. *Journal of Colloid and Interface Science* 111:466-474.
63. Berzofsky, J.A., I.J. Berkower, K.B. Cease, G.K. Buckenmeyer, H.Z. Streicher, and C. Delisi. 1986. Structural and conformational requirements for protein antigen recognition by MHC class II - restricted T cells and clones. In *Modern Approaches to Vaccines*. F. Brown, R. Chanock, and R. Lerner, editors. Cold Spring Harbor Laboratory, New York. 123-127.
64. Berzofsky, J.A., K.B. Cease, G.K. Buckenmeyer, H.Z. Streicher, C. Delisi, and I.J. Berkower. 1986. Structural and conformational requirements for Ir-gene-controlled myoglobin epitope recognition by Ia-restricted T cells and clones. In *Regulation of Immune Gene Expression*. M. Feldmann, and A. McMichael, editors. Humana Press, Clifton, NJ. 225-234.
65. Kawamura, H., and J.A. Berzofsky. 1986. Enhancement of antigenic potency in vitro and immunogenicity in vivo by coupling the antigen to anti-immunoglobulin. *J. Immunol.* 136:58-65.
66. Streicher, H.Z., F. Cuttitta, G.K. Buckenmeyer, H. Kawamura, J. Minna, and J.A. Berzofsky. 1986. Mapping the idiotopes of a monoclonal anti-idiotypic antibodies: Detection of a common idiotope. *J. Immunol.* 136:1007-1014.
67. Berzofsky, J.A. 1987. *Ir* genes: antigen-specific genetic regulation of the immune response. In *The Antigens*. M. Sela, editor. Academic Press, New York. 1-146.
68. Berkower, I., G.K. Buckenmeyer, and J.A. Berzofsky. 1986. Molecular mapping of a histocompatibility - restricted immunodominant T cell epitope with synthetic and natural peptides: Implications for antigenic structure. *J. Immunol.* 136:2498-2503.

69. Berzofsky, J.A. 1986. Structural features of protein antigenic sites recognized by helper T cells: What makes a site immunodominant?. In *The Year in Immunology 1985-1986*. J.M. Cruse, and R.E. Lewis, Jr., editors. Karger, Basel. 28-38.
70. Simpson, E., R. Lieberman, I. Ando, D.H. Sachs, W.E. Paul, and J.A. Berzofsky. 1986. How many class II Immune Response Genes? A reappraisal of the evidence. *Immunogenetics* 23:302-308.
71. Delisi, C., J. Cornette, H. Margalit, K. Cease, J. Spouge, and J.A. Berzofsky. 1987. The role of amphipathicity as an indicator of T cell antigenic sites on proteins. In *Immunogenicity of Protein Antigens: Repertoire and Regulation*. E. Sercarz, and J.A. Berzofsky, editors. CRC Press, Boca Raton. 35-42.
72. Berzofsky, J.A., J. Cornette, H. Margalit, I. Berkower, K. Cease, and C. Delisi. 1986. Molecular features of class II MHC-restricted T cell recognition of protein and peptide antigens: the importance of amphipathic structures. In *Current Topics in Microbiology and Immunology* Vol. 130. F. Melchers, and H. Koprowski, editors. Springer-Verlag, New York. 14-24.
73. Kawamura, H., S.O. Sharrow, D.W. Alling, D. Stephany, J. York-Jolley, and J.A. Berzofsky. 1986. IL-2 receptor expression in unstimulated murine splenic T cells: Localization to L3T4+ cells and regulation by non-H-2-linked genes. *J. Exp. Med.* 86:1376-1390.
74. Wagner, D.K., J. York-Jolley, T.R. Malek, J.A. Berzofsky, and D.L. Nelson. 1986. Antigen-specific murine T cell clones produce soluble interleukin 2 receptor on stimulation with specific antigens. *J. Immunol.* 137:592-596.
75. Good, M.F., J.A. Berzofsky, W.L. Maloy, Y. Hayashi, N. Fujii, W.T. Hockmeyer, and L.H. Miller. 1986. Genetic control of the immune response in mice to a Plasmodium faciparum sporozoite vaccine: Widespread non-responsiveness to a single malaria T epitope in highly repetitive vaccine. *J. Exp. Med.* 164:655-660.
76. Berzofsky, J.A., K.B. Cease, I.J. Berkower, H. Margalit, J. Cornette, J. Spouge, C. Spencer, G. Buckenmeyer, H. Streicher, M. Kojima, and C. Delisi. 1988. The role of MHC and amphipathic structures in T cell recognition: Features determining immunodominance. In *Antigen Processing and Presentation*. H. Vogel, B. Pernis, and S. Silverstein, editors. Columbia University Press, New York. 125-131.
77. Finnegan, A., M.A. Smith, J.A. Smith, J.A. Berzofsky, D.H. Sachs, and R.J. Hodes. 1986. The T cell repertoire for recognition of a phylogenetically distant protein antigen: peptide specificity and MHC restriction of staphylococcal nuclease specific T cell clones. *J. Exp. Med.* 164:897-910.
78. Kawamura, H., and J.A. Berzofsky. 1987. Influence of MHC-linked genes on idotype expression. In *Immunogenicity of Protein Antigens: Repertoire and Regulation* Vol.2. E.E. Sercarz, and J.A. Berzofsky, editors. CRC Press, Boca Raton. 121-127.

79. Berzofsky, J.A., K.B. Cease, I.J. Berkower, H. Margalit, J. Cornette, and C. Delisi. 1986. Immunodominance of amphipathic peptides and their localization on the cell surface for antigen presentation to helper T cells. In *Progress in Immunology V*. B. Cinader, and R.G. Miller, editors. Academic Press, New York. 255-265.
80. Cease, K.B., I. Berkower, J. York-Jolley, and J.A. Berzofsky. 1986. T cell clones specific for an amphipathic alpha helical region of sperm whale myoglobin show differing fine specificities for synthetic peptides: A multi-view/single structure interpretation of immunodominance. *J. Exp. Med.* 164:1779-1784.
81. Cease, K.B., G. Buckenmeyer, I. Berkower, J. York-Jolley, and J.A. Berzofsky. 1986. Immunologically relevant peptide antigen exists on the presenting cell in a manner accessible to macromolecules in solution. *J. Exp. Med.* 164:1440-1454.
82. Berzofsky, J.A., and K.B. Cease. 1986. Peptide antigenic structure and localization on the surface of the presenting cell for antigen presentation. *Annales de l'Institut Pasteur* 137D:312-316.
83. Berzofsky, J.A., K.B. Cease, S. Ozaki, H. Margalit, J.L. Cornette, J. Spouge, and C. Delisi. 1987. Helper T cell immunity: Implications for vaccines. In *Vaccines '87*. R. Chanock, F. Brown, H. Ginsberg, and R. Lerner, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. 26-32.
84. Finnegan, A., and J.A. Berzofsky. 1986. Antigen recognition by T cells. *Immunol. Today* 7:317-319.
85. Spouge, J.L., H.R. Guy, J.L. Cornette, H. Margalit, K. Cease, J.A. Berzofsky, and C. Delisi. 1987. Strong conformational propensities enhance T-cell antigenicity. *J. Immunol.* 138:204-212.
86. Good, M.F., I.A. Quakyi, A. Saul, J.A. Berzofsky, R. Carter, and L.H. Miller. 1987. Human T clones reactive to the sexual stages of *Plasmodium falciparum* malaria: High frequency of gamete-reactive T cells in peripheral blood of non-exposed donors. *J. Immunol.* 138:306-311.
87. Coico, R.F., J.A. Berzofsky, J. York-Jolley, S. Ozaki, G.W. Siskind, and G.J. Thorbecke. 1987. Physiology of IgD. VII. Induction of receptors for IgD on cloned T cells by IgD and Interleukin 2. *J. Immunol.* 138:4-6.
88. Ozaki, S., J. York-Jolley, H. Kawamura, and J.A. Berzofsky. 1987. Cloned protein antigen-specific, Ia-restricted T cells with both helper and cytolytic activities: Mechanisms of activation and killing. *Cellular Immunol.* 105:301-316.
89. Good, M.F., J.A. Berzofsky, L. Maloy, M.N. Lunde, Y. Hayashi, N. Fujii, W.T. Hockmeyer, B. Moss, and L.H. Miller. 1987. Genetic control of the immune response to a *Plasmodium falciparum* sporozoite vaccine and to the Circumsporozoite (CS)

- protein. In *Vaccines '87*. R. Chanock, F. Brown, H. Ginsberg, and R. Lerner, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York . 81-85.
90. Margalit, H., J.L. Spouge, J.L. Cornette, K. Cease, C. Delisi, and J.A. Berzofsky. 1987. Prediction of immunodominant helper T-cell antigenic sites from the primary sequence. *J. Immunol.* 138:2213-2229.
 91. Takacs, L., J.A. Berzofsky, J. York-Jolley, T. Akahoshi, E. Blasi, and S.K. Durum. 1987. IL 1-Induction by murine T cell clones: Detection of an IL 1-inducing lymphokine. *J. Immunol.* 138:2124-2131.
 92. Good, M.F., W.L. Maloy, M.N. Lunde, H. Margalit, J.L. Cornette, G.L. Smith, B. Moss, L.H. Miller, and J.A. Berzofsky. 1987. Construction of a synthetic immunogen: use of a new T-helper epitope on malaria circumsporozoite protein. *Science* 235:1059-1062.
 93. Hoffman, S.L., L.T. Cannon, J.A. Berzofsky, W.R. Majarian, J.G. Young, W.L. Maloy, and W.T. Hockmeyer. 1987. Plasmodium falciparum: Increased immunity with T cell epitope on sporozoite vaccine. *Exp. Parasitology* 64:64-70.
 94. Cornette, J.L., K.B. Cease, H. Margalit, J.L. Spouge, J.A. Berzofsky, and C. Delisi. 1987. Hydrophobicity scales and computational techniques for detecting amphipathic structures in proteins. *J. Mol. Biol.* 195:659-686.
 95. Cease, K.B., H. Margalit, J.L. Cornette, S.D. Putney, W.G. Robey, C. Ouyang, H.Z. Streicher, P.J. Fischinger, R.C. Gallo, C. Delisi, and J.A. Berzofsky. 1987. Helper T cell antigenic site identification in the AIDS virus gp120 envelope protein and induction of immunity in mice to the native protein using a 16-residue synthetic peptide. *Proc. Natl. Acad. Sci. USA* 84:4249-4253.
 96. Ozaki, S., and J.A. Berzofsky. 1987. Antibody conjugates mimic specific B cell presentation of antigen: Relationship between T and B cell specificity. *J. Immunol.* 138:4133-4142.
 97. Berzofsky, J.A., K.B. Cease, J.L. Cornette, J.L. Spouge, H. Margalit, I.J. Berkower, M.F. Good, L.H. Miller, and C. Delisi. 1987. Protein antigenic structures recognized by T cells: Potential applications to vaccine design. *Immunol. Rev.* 98:9-52.
 98. Berzofsky, J.A. 1988. Features of T-cell recognition and antigen structure useful in the design of vaccines to elicit T-cell immunity. *Vaccines* 6:89-93.
 99. Berzofsky, J.A., and S. Ozaki. 1988. Antibody conjugates mimic specific B-cell presentation of antigen: epitope restrictions in specific T-B interaction. In *Antigen Presenting Cells: Diversity, Differentiation and Regulation*. L.B. Schook, and J.G. Tew, editors. Alan R. Liss, Inc., New York. 41-48.
 100. Good, M.F., D. Pombo, I.A. Quakyi, E.M. Riley, R.A. Houghten, A. Menon, D.W. Alling, J.A. Berzofsky, and L.H. Miller. 1988. Human T cell recognition of the

- circumsporozoite protein of *Plasmodium falciparum*. Immunodominant T cell domains map to the polymorphic regions of the molecule. *Proc. Natl. Acad. Sci. USA* 85:1199-1203.
101. Good, M.F., J.A. Berzofsky, and L.H. Miller. 1988. The T-cell response to the malaria circumsporozoite protein: an immunological approach to vaccine development. *Annu. Rev. Immunol.* 6:663-688.
 102. Lin, J., J.A. Berzofsky, and T.L. Delovitch. 1988. Ultrastructural study of internalization and recycling of antigen by antigen presenting cells. *J. Mol. Cell. Immunol.* 3:321-343.
 103. Kojima, M., K.B. Cease, G.K. Buckenmeyer, and J.A. Berzofsky. 1988. Limiting dilution comparison of the repertoires of high and low responder MHC-restricted T cells. *J. Exp. Med.* 167:1100-1113.
 104. Good, M.F., D. Pombo, W.L. Maloy, V.F. De la Cruz, L.H. Miller, and J.A. Berzofsky. 1988. Parasite polymorphism present within minimal T-cell epitopes of *Plasmodium falciparum* circumsporozoite protein. *J. Immunol.* 140:1645-1650.
 105. Takahashi, H., J. Cohen, A. Hosmalin, K.B. Cease, R. Houghten, J. Cornette, C. Delisi, B. Moss, R.N. Germain, and J.A. Berzofsky. 1988. An immunodominant epitope of the HIV gp160 envelope glycoprotein recognized by class I MHC molecule-restricted murine cytotoxic T lymphocytes. *Proc. Natl. Acad. Sci. USA* 85:3105-3109.
 106. Darst, S.A., C.R. Robertson, and J.A. Berzofsky. 1988. Adsorption of the protein antigen myoglobin affects the binding of conformation specific monoclonal antibodies. *Biophysical Journal* 53:533-539.
 107. Pombo, D., W.L. Maloy, J.A. Berzofsky, and M.F. Good. 1988. Neonatal exposure to immunogenic peptides. Differential susceptibility to tolerance induction of helper T cells and B cells reactive to malarial circumsporozoite peptide epitopes. *J. Immunol.* 140:3594-3598.
 108. Berzofsky, J.A. 1988. T-cell recognition of protein and peptide antigens: applications to vaccines development. In *Technological Advances in Vaccine Development*. L. Laskey, editor. Alan R. Liss, Inc., New York, 587-602.
 109. Berzofsky, J.A. 1988. Immunodominance in T lymphocyte recognition. *Immunol. Letters* 18:83-92.
 110. Ozaki, S., S.K. Durum, K. Muegge, J. York-Jolley, and J.A. Berzofsky. 1988. Production of T-T hybrids from T cell clones: Direct comparison between cloned T cells and T hybridoma cells derived from them. *J. Immunol.* 141:71-78.
 111. Brett, S.J., K.B. Cease, and J.A. Berzofsky. 1988. Influences of antigen processing on the expression of the T cell repertoire: Evidence for MHC-specific hindering structures on the products of processing. *J. Exp. Med.* 168:357-373.

112. Good, M.F., D. Pombo, M.N. Lunde, W.L. Maloy, R. Halenbeck, K. Koths, L.H. Miller, and J.A. Berzofsky. 1988. Recombinant human interleukin-2 (IL-2) overcomes genetic nonresponsiveness to malaria sporozoite peptides. Correlation of effect with biological activity of IL-2. *J. Immunol.* 141:972-977.
113. Cease, K.B., and J.A. Berzofsky. 1990. T cell immunity and vaccine engineering: Application to the AIDS virus. In *AIDS Vaccine Research and Clinical Trials*. S. D. Putney, and D. P. Bolognesi, editors. Marcell Dekker, New York, pp. 139-156.
114. Kumar, S., L.H. Miller, I.A. Quakyi, D.B. Keister, R.A. Houghten, W.L. Maloy, B. Moss, J.A. Berzofsky, and M.F. Good. 1988. Cytotoxic T cells specific for the circumsporozoite protein of *Plasmodium falciparum*. *Nature* 334:258-260.
115. Berzofsky, J.A., A. Bensussan, K.B. Cease, J.F. Bourge, R. Cheynier, Z. Lurhuma, J.-J. Salaün, R.C. Gallo, G.M. Shearer, and D. Zagury. 1988. Antigenic peptides recognized by T lymphocytes from AIDS viral envelope-immune humans. *Nature* 334:706-708.
116. Margalit, H., C. Delisi, and J.A. Berzofsky. 1990. Computer predictions of T-cell epitopes. In *New Generation Vaccines*. G.C. Woodrow, and M.M. Levine, editors. Marcel Dekker, New York, 109-116.
117. Berzofsky, J.A. 1989. Structural features of T-cell recognition: Applications to vaccine design. *Phil. Trans. Roy. Soc. London B.* , 323: 535-544.
118. Cease, K.B., and J.A. Berzofsky. 1988. Antigenic structures recognized by T cells: Toward the rational design of an AIDS vaccine. *AIDS* 2:(suppl.1) S95-S101.
119. Berzofsky, J.A., S.J. Brett, H.Z. Streicher, and H. Takahashi. 1988. Antigen processing for presentation to T lymphocytes: Function, mechanisms, and implications for the T-cell repertoire. *Immunol. Rev.* 106: 5-31.
120. Good, M.F., L.H. Miller, S. Kumar, I.A. Quakyi, D. Keister, J. Adams, B. Moss, J.A. Berzofsky, and R. Carter. 1988. Limited immunological recognition of critical malaria vaccine candidate antigens. *Science* 242: 574-577.
121. Berzofsky, J.A., and I.J. Berkower. 1989. Immunogenicity and antigen structure. In *Fundamental Immunology* 2nd Edition. W.E. Paul, editor. Raven Press, New York, pp. 169-208.
122. Berzofsky, J.A., S.L. Epstein, and I.J. Berkower. 1989. Antigen-antibody interactions and monoclonal antibodies. In *Fundamental Immunology* 2nd edition. W.E. Paul, editor. Raven Press, New York, pp. 315-356.
123. Dontfraid, F., M.A. Cochran, D. Pombo, J.D. Knell, I. Quakyi, S. Kumar, R.A. Houghten, J.A. Berzofsky, L.H. Miller, and M.F. Good. 1988. Human and murine CD4 T-cell epitopes map to the same region of the malaria circumsporozoite protein:

- Limited immunogenicity of sporozoites and circumsporozoite protein. *Mol. Biol. Med.*, 5: 185-196.
124. Hosmalin, A.M., P.L. Nara, M. Zweig, K.B. Cease, E. Gard, P.D. Markham, S.D. Putney, M.D. Daniel, R.C. Desrosiers, and J.A. Berzofsky. 1989. Enhancement of an antibody response to the envelope glycoprotein of HIV-1 in primates by priming with helper T-cell epitope peptides. In *Vaccines 89*. R. Lerner, H. Ginsberg, R. Chanock, and F. Brown, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, 121-124.
 125. Takahashi, H., J. Cohen, A. Hosmalin, K.B. Cease, R. Houghten, J.L. Cornette, C. Delisi, S. Merli, B. Moss, R.N. Germain, and J.A. Berzofsky. 1989. Limited epitope repertoire recognized with class I MHC molecules by murine cytotoxic lymphocytes on the HIV gp160 envelope glycoprotein. In *Vaccines 89*. R. Lerner, H. Ginsberg, R. Chanock, and F. Brown, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor., 109-114.
 126. Berzofsky, J.A. 1989. Immunodominance of T-cell epitopes: Applications to vaccine design. In *Vaccines 89*. R. Lerner, H. Ginsberg, R. Chanock, and F. Brown, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, 27-31.
 127. Berzofsky, J.A. 1988. The structural basis of antigen recognition by T lymphocytes: Implications for vaccines. *J. Clin. Invest.* 82: 1811-1817.
 128. Berzofsky, J.A. 1989. Mechanisms of immunodominance in T-cell recognition, with applications to vaccine design. In *Immune System and Cancer: Proceedings of the Nineteenth International Symposium of the Princess Takamatsu Cancer Research Fund*. T. Hamaoka, R.J. Hodes, G. Klein, T. Sugimura, S. Takayama, and Y. Yamamura, editors. The Princess Takamatsu Cancer Research Fund, Tokyo, 161-177.
 129. Takahashi, H., K.B. Cease, and J. A. Berzofsky. 1989. Identification of proteases that process distinct epitopes on the same protein. *J. Immunol.* 142: 2221-2229.
 130. Brett, S. J., D. McKean, J. York-Jolley, and J. A. Berzofsky. 1989. Antigen presentation to specific T cells by Ia molecules selectively altered by site-directed mutagenesis. *Internat. Immunol.* 1: 130-140.
 131. Berzofsky, J. A., and Y. Paterson. 1990. Discussion summary: Antigen structure: B cell and T cell determinants and immunogenicity. In *Immunogenicity (UCLA Symposia on Molecular and Cellular Biology, New Series, Vol. 113)*. C. Janeway, J. Sprent, and E. Sercarz, editors. Alan R. Liss, New York, pp. 65-67.
 132. Hoffman, S. L., J. A. Berzofsky, D. Isenbarger, E. Zeltzer, W. R. Majarian, M. Gross, and W. R. Ballou. 1989. Immune response gene regulation of immunity to *Plasmodium berghei* sporozoites and circumsporozoite protein vaccines: Overcoming genetic restriction with whole organism and subunit vaccines. *J. Immunol.* 142: 3581-3584.

133. Cornette, J. L., H. Margalit, C. DeLisi, and J. A. Berzofsky. 1989. Concepts and methods in the identification of T cell epitopes and their use in the construction of synthetic vaccines. *Methods in Enzymol.* 178: 611-634.
134. Sanui, H., T. M. Redmond, S. Kotake, B. Wiggert, L.-H. Hu, H. Margalit, J. A. Berzofsky, G. J. Chader, and I. Gery. 1989. Identification of an immunodominant and highly immunopathogenic determinant in the retinal interphotoreceptor retinoid-binding protein (IRBP). *J. Exp. Med.* 169: 1947-1960.
135. DeGroot, A. S., A. H. Johnson, W. L. Maloy, I. A. Quakyi, E. M. Riley, A. Menon, S. M. Banks, J. A. Berzofsky, and M. F. Good. 1989. Human T cell recognition of polymorphic epitopes from malaria circumsporozoite protein. *J. Immunol.* 142: 4000-4005.
136. Palker, T. J., T. J. Matthews, A. Langlois, M. E. Tanner, M. E. Martin, R.M. Searce, J. E. Kim, J. A. Berzofsky, D. P. Bolognesi, and B. F. Haynes. 1989. Polyvalent human immunodeficiency virus synthetic immunogen comprised of envelope gp120 T helper cell sites and B cell neutralization epitopes. *J. Immunol.* 142: 3612-3619.
137. Sanza, L. R., L.M.Gierasch, J. A. Berzofsky, G. K. Buckenmeyer, K. B. Cease, and C. S. Ouyang. 1988. Formation of amphipathic secondary structure is correlated to T-cell antigenicity in a series of synthetic peptides from sperm whale myoglobin. In *Peptides: Chemistry and Biology*. G. R. Marshall, editor, ESCOM, Leiden, pp. 549-550.
138. Palker, T. J., T. J. Matthews, A. Langlois, J. E. Kim, J. A. Berzofsky, D. P. Bolognesi, and B. F. Haynes. 1989. Multivalent synthetic peptide inoculum induces neutralizing antibodies to divergent HIV isolates and anamnestic T-cell responses to HIV gp120. In *Early Human Retroviruses*. J. E. Groopman, I. Chen, M. Essex, and R. Weiss, editors. Alan R. Liss, New York, in press.
139. Clerici, M., N. I. Stocks, R. A. Zajac, R. N. Boswell, D. C. Bernstein, D. L. Mann, G. M. Shearer, and J. A. Berzofsky. 1989. IL-2 production used to detect antigenic peptide recognition by T helper lymphocytes from asymptomatic, HIV seropositive individuals. *Nature* 339: 383-385.
140. Williams, W.V., S.D. London, D.B. Weiner, S. Wadsworth, J.A. Berzofsky, F. Robey, D.H. Rubin, and M.I. Greene. 1989. Immune response to a molecularly defined internal image idiotope. *J. Immunol.* 142: 4392-4400.
141. Brett, S.J., K.B. Cease, C.S. Ouyang, and J.A. Berzofsky. 1989. Fine specificity of T cell recognition of the same peptide in association with different I-A molecules. *J. Immunol.* 143: 771-779.
142. Hale, P.M., K.B. Cease, R.A. Houghten, C. Ouyang, S. Putney, K. Javaherian, H. Margalit, J.L. Cornette, J.L. Spouge, C. Delisi, and J.A. Berzofsky. 1989. T cell multideterminant regions in the human immunodeficiency virus envelope: toward

overcoming the problem of major histocompatibility complex restriction. *Internat. Immunol.* 1: 409-415.

143. Berzofsky, J.A., A. Kurata, H. Takahashi, S.J. Brett, and D.J. McKean. 1989. Molecular studies of antigen processing and presentation to T cells by class II MHC molecules. In Cold Spring Harbor Symposium on Quantitative Biology LIV: Immunological Recognition. J.D. Watson, and J. R. Inglis, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, pp. 417-430.
144. Kurata, A., T.J. Palker, R.D. Streilein, R.M. Searce, B.F. Haynes, and J.A. Berzofsky. 1989. Immunodominant sites of human T-cell lymphotropic virus type 1 envelope protein for murine helper T cells. *J. Immunol.* 143: 2024-2030.

145. Redmond, T.M., H. Sanui, L.H. Hu, B. Wiggert, H. Margalit, J.A. Berzofsky, G.J. Chader, and I. Gery. 1989. Immune responses to peptides derived from the retinal protein IRBP: immunopathogenic determinants are not necessarily immunodominant. *Clinical Immunology and Immunopathology* 53: 212-224.
146. Lark, L.R., J.A. Berzofsky, and L.M. Gierasch. 1989. T-Cell antigenic peptides from sperm whale myoglobin fold as amphipathic helices: a possible determinant for immunodominance?. *Peptide Research* : 2: 314-321.
147. Vacchio, M.S., J.A. Berzofsky, U. Krzych, J.A. Smith, R.J. Hodes, and A. Finnegan. 1989. Sequences outside a minimal immunodominant site exert negative effects on recognition by staphylococcal nuclease-specific T-cell clones. *J. Immunol.* 143: 2814-2819.
148. Vajda, S., H. Margalit, R. Kataoka, J. L. Cornette, J. A. Berzofsky, and C. DeLisi. 1990. Molecular structure and vaccine design. *Ann. Rev. Biophys. Biophys. Chem.* 19: 69-82..
149. Takahashi, H., S. Merli, S.D. Putney, R. Houghten, B. Moss, R.N. Germain, and J.A. Berzofsky. 1989. A single amino acid interchange yields reciprocal CTL specificities for HIV gp160. *Science* 246: 118-121.
150. Takahashi, H., R. Houghten, S.D. Putney, D.H. Margulies, B. Moss, R.N. Germain, and J.A. Berzofsky. 1989. Structural requirements for class-I MHC molecule-mediated antigen presentation and cytotoxic T-cell recognition of an immunodominant determinant of the HIV envelope protein. *J. Exp. Med.* 170: 2023-2035.
151. Weiss, W.R., M.F. Good, M.R. Hollingdale, L.H. Miller, and J.A. Berzofsky. 1989. Genetic control of immunity to *Plasmodium yoelii* sporozoites. *J. Immunol.* 143: 4263-4266.
152. Berzofsky, J.A., P.M. Hale, M. Clerici, K.B. Cease, R.A. Houghten, S.D. Putney, R.A. Zajac, R.N. Boswell, H. Margalit, J.L. Cornette, J.L. Spouge, C. Delisi, and G.M. Shearer. 1990. Multideterminant regions of the HIV-1 envelope with sites seen by murine and human helper T cells: circumventing the MHC restriction problem. In *Vaccines 90*. F. Brown, R.M. Chanock, H.S. Ginsberg, and R.A. Lerner, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, pp. 307-312.
153. Takahashi, H., R. Houghten, S. Merli, S.D. Putney, B. Moss, R.N. Germain, and J.A. Berzofsky. 1990. Immunodominant CTL epitope of HIV-1 envelope protein: the relationship between viral mutation and CTL specificity. In *Vaccines 90*. F. Brown, R.M. Chanock, H.S. Ginsberg, and R.A. Lerner, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, pp 269-276.
154. Vinetz, J.M., S. Kumar, M.F. Good, B.J. Fowlkes, J.A. Berzofsky, and L.H. Miller. 1990. Adoptive transfer of CD8+ T cells from immune animals does not transfer immunity to blood stage *Plasmodium yoelii* malaria. *J. Immunol.* 144: 1069-1074.

155. Takahashi, H., R.N. Germain, B. Moss, and J.A. Berzofsky. 1990. An immunodominant class I-restricted CTL determinant of HIV-1 induces CD4 class II-restricted help for itself. *J. Exp. Med.* 171: 571-576.
156. Weiss, W.R., S. Mellouk, R.A. Houghten, M. Sedegah, S. Kumar, M.F. Good, J.A. Berzofsky, L.H. Miller, and S.L. Hoffman. 1990. Cytotoxic T cells recognize a peptide from the circumsporozoite protein on malaria infected hepatocytes. *J. Exp. Med.* 171: 763-773.
157. Kotake, S., B. Wiggert, T.M. Redmond, D.E. Borst, J.M. Nickerson, H. Margalit, J.A. Berzofsky, G.J. Chader, and I. Gery. 1990. Repeated determinants within the retinal interphotoreceptor retinoid-binding protein (IRBP). Immunological properties of the repeats of an immunodominant determinant. *Cellular Immunol.*, 126: 331-342.
158. Hosmalin, A., M. Clerici, R. Houghten, C.D. Pendleton, C. Flexner, D.R. Lucey, B. Moss, R.N. Germain, G.M. Shearer, and J.A. Berzofsky. 1990. An epitope in HIV-1 reverse transcriptase recognized by both mouse and human CTL. *Proc. Natl. Acad. Sci. U. S. A.* 87: 2344-2348.
159. Ahlers, J., M. Clerici, A. Hosmalin, G.M. Shearer, and J.A. Berzofsky. 1990. Host immune response: T helper cell responses. In *Techniques in HIV Research*. A. Aldovini, and B. Walker, editors. Stockton Press, New York, pp. 211-222.
160. Berzofsky, J.A. 1990. Immunogenicity of antigens recognized by T cells. In *Cellular Immunity and the Immunotherapy of Cancer*. M.T. Lotze, and O.J. Finn, editors. Wiley-Liss, Inc., New York, 1-22.
161. Takahashi, H., T. Takeshita, B. Morein, S.D. Putney, R.N. Germain, and J.A. Berzofsky. 1990. Induction of specific CD8⁺ class I-MHC restricted CTL against HIV envelope protein by immunization with purified whole protein in ISCOMs. *Nature*, 344: 873-875.
162. Kim, J.E., M. Kojima, R. Houghten, C.D. Pendleton, J.L. Cornette, C. Delisi, and J.A. Berzofsky. 1990. Characterization of a helper T-cell epitope recognized by mice of a low responder major histocompatibility type. *Molec. Immunol.* 27: 941-946.
163. Weiner, D.B., W.V. William, M.J. Merva, K. Huebner, J.A. Berzofsky, and M.I. Greene. 1990. HIV Infectivity analysis of viral envelope determinants and target cell requirements for infectivity by HIV-1. In *Vaccines 90*. F. Brown, R.M. Chanock, H.S. Ginsberg, and R.A. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. 339-345.
164. Kurata, A., and J.A. Berzofsky. 1990. Analysis of peptide residues interacting with MHC molecule or T-cell receptor: can a peptide bind in more than one way to the same MHC molecule?. *J. Immunol.* 144: 4526-4535.

165. Kumar, S., J. Gorden, J.L. Flynn, J.A. Berzofsky, and L.H. Miller. 1990. Immunization of mice against *Plasmodium vinckei* with a combination of attenuated *Salmonella typhimurium* and malarial antigen. *Infect. Immun.* 58: 3425-3429.
166. Kozlowski, S., T. Takeshita, W. -H. Boehncke, H. Takahashi, L. F. Boyd, R. N. Germain, J. A. Berzofsky, and D. H. Margulies. 1991. Excess β 2-microglobulin promotes functional peptide association with purified soluble class I MHC molecules. *Nature* 349: 74-77.
167. Hosmalin, A., P. L. Nara, M. Zweig, N. W. Lerche, K. B. Cease, E. A. Gard, P. D. Markham, S. D. Putney, M. D. Daniel, R. C. Desrosiers, and J. A. Berzofsky. 1991. Priming with helper T-cell epitope peptides enhances the antibody response to the envelope glycoprotein of HIV 1 in primates. *J. Immunol.* 146: 1667-1673.
168. Berzofsky, J.A. 1991. Progress towards an artificial vaccine for HIV: identification of helper and cytotoxic T-cell epitopes and methods of immunization. *Biotech. Therapeutics.* 2: 123-135.
169. Berzofsky, J.A. 1991. Approaches and issues in the development of vaccines against HIV. *J. Acq. Immune Defic. Syndromes* 4: 451-459.
170. Clerici, M., D. R. Lucey, R. A. Zajac, R. N. Boswell, H. M. Gebel, H. Takahashi, J. A. Berzofsky, and G. M. Shearer. 1991. Detection of cytotoxic T lymphocytes specific for synthetic peptides of gp160 in HIV-seropositive individuals. *J. Immunol.* 146: 2214-2219.
171. Culver, K.W., C. Able, R. Toper, S. Freeman, J. A. Berzofsky, and R. M. Blaese. 1990. Lymphocytes as vehicles for gene therapy. In *Cellular Immunity and the Immunotherapy of Cancer*. M.T. Lotze and O.J. Finn, editors. Wiley-Liss, Inc., New York. 129-137.
172. Clerici, M., C. O. Tacket, C. S. Via, D. R. Lucey, S. C. Muluk, R. A. Zajac, R. N. Boswell, J. A. Berzofsky, and G. M. Shearer. 1991. Immunization with subunit human immunodeficiency virus vaccine generates stronger T helper cell immunity than natural infection. *Eur. J. Immunol.* 21: 1345-1349.
173. Takahashi, H., T. Takeshita, B. Moreln, S. Putney, R. N. Germain, and J. A. Berzofsky. 1991. A unique subunit immunogen, ISCOM-gp160, can elicit MHC class-I-restricted HIV envelope-specific CD8⁺CTLs. In *Vaccines 91*. R.M. Chanock, H.S. Ginsberg, F. Brown, and R.A. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. 1-7.
174. Lipham, W.J., T. M. Redmond, H. Takahashi, J. A. Berzofsky, B. Wiggert, G. J. Chader, and I. Gery. 1991. Recognition of peptides that are immunopathogenic but cryptic: Mechanisms that allow lymphocytes sensitized against cryptic peptides to initiate pathogenic autoimmune processes. *J. Immunol.* 146: 3757-3762.

175. Clerici, M., J. A. Berzofsky, G. M. Shearer, and C. O. Tacket. 1991. Exposure to HIV-1 indicated by HIV-specific T helper cell responses before detection of infection by polymerase chain reaction and serum antibodies. *J. Infect. Dis.* 164: 178-182.
176. Berzofsky, J.A. 1991. Mechanisms of T cell recognition with application to vaccine design. *Molec. Immunol.* 28:217-223.
177. Berzofsky, J.A. 1991. Development of artificial vaccines against HIV using defined epitopes. *FASEB J.* 5: 2412-2418.
178. Berzofsky, J.A. 1991. Antigenic peptide interaction with MHC molecules: implications for the design of artificial vaccines. *Seminars in Immunology* 3: 203-216.
179. Goodman-Snitkoff, G., M. F. Good, J. A. Berzofsky, and R. J. Mannino. 1991. Role of intrastructural/intermolecular help in immunization with peptide-phospholipid complexes. *J. Immunol.* 147: 410-415.
180. Berzofsky, J.A., C. D. Pendleton, M. Clerici, J. Ahlers, D. R. Lucey, S. D. Putney, and G. M. Shearer. 1991. Construction of peptides encompassing multideterminant clusters of human immunodeficiency virus envelope to induce in vitro T-cell responses in mice and humans of multiple MHC types. *J. Clin. Invest.* 88: 876-884.
181. Berzofsky, J.A. 1991. Human immunodeficiency virus: structural features of T-cell epitopes and their use in vaccine development. In *Viruses and the Cellular Immune Response*. D.B. Thomas, editor. Marcel Dekker, Inc., New York.
182. Berzofsky, J.A., C. D. Pendleton, M. Clerici, J. Ahlers, D. R. Lucey, S. D. Putney, and G. M. Shearer. 1991. Peptides containing multideterminant clusters of Human Immunodeficiency Virus envelope induce murine and human T -cell responses in diverse histocompatibility types. *Transac. Assoc. Amer. Phys.* 104: 69-77.
183. Ida, H., A. Kurata, K. Eguchi, A. Kawakami, K. Migita, T. Fukuda, T. Nakamura, Y. Kusumoto, J. A. Berzofsky, and S. Nagataki. 1991. Different B-cell responses to human T-cell lymphotropic virus type I (HTLV-I) envelope synthetic peptides in HTLV-I-infected individuals. *Journal of Clinical Immunology* 11:143-151.
184. DeGroot, A. S., M. Clerici, A. Hosmalin, S. H. Hughes, D. Barnd, C. W. Hendrix, R. A. Houghten, G. M. Shearer, and J. A. Berzofsky. 1991. Human immunodeficiency virus (HIV-1) reverse transcriptase T helper epitopes identified in mice and humans: Correlation with a cytotoxic T cell (CTL) epitope. *J. Infect. Dis.* 164: 1058-1065.
185. Takahashi, H., Y. Nakagawa, C. D. Pendleton, R. A. Houghten, K. Yokomuro, R. N. Germain, and J. A. Berzofsky. 1992. Analysis of CTL crossreactivity to an HIV-1 immunodominant determinant: elicitation of widely crossreactive CTL. In *Vaccines 92*. F. Brown, R. Chanock, H. Ginsberg, and R. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. 69-74.

186. Shirai, M. and J. A. Berzofsky. 1992. Promiscuity of cytotoxic T-cell epitopes from the HIV-1 envelope . In *Vaccines 92*. F. Brown, R. Chanock, H. Ginsberg, and R. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. 243-246.
187. Berzofsky, J. A. 1992. Progress towards artificial vaccines for HIV. In *Vaccines 92*. F. Brown, R. Chanock, H. Ginsberg, and R. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. 41-50.
188. Takahashi, H., Y. Nakagawa, C. D. Pendleton, R. A. Houghten, K. Yokomuro, R. N. Germain, and J. A. Berzofsky. 1992. Induction of broadly crossreactive cytotoxic T cells recognizing an HIV-1 envelope determinant. *Science* , 255: 333-336.
189. Shirai, M., C. D. Pendleton, and J. A. Berzofsky. 1992. Broad recognition of cytotoxic T-cell epitopes from the HIV-1 envelope protein with multiple class I histocompatibility molecules. *J. Immunol.* 148: 1657-1667.
190. Cornette, J. L., H. Margalit, C. DeLisi, and J. A. Berzofsky. 1993. The amphipathic helix as a structural feature involved in T-cell recognition. In *The Amphipathic Helix*. R.M. Epand, editor. CRC Press, Boca Raton., 333-346.
191. Clerici, M., J. V. Giorgi, C. -C. Chou, V. K. Gudeman, J. A. Zack, P. Gupta, H. -N. Ho, P. G. Nishanian, J. A. Berzofsky, and G. M. Shearer. 1992. Cell mediated immune response to human immunodeficiency virus type I (HIV-1) in seronegative homosexual men with recent sexual exposure to HIV-1. *J. Infect. Dis.* 165: 1012-1019.
192. Kozlowski, S., M. Corr, T. Takeshita, L. F. Boyd, C. D. Pendleton, R. N. Germain, J. A. Berzofsky, and D. H. Margulies. 1992. Serum angiotensin-1 converting enzyme activity processes an HIV 1 gp160 peptide for presentation by MHC class I molecules. *J. Exp. Med.* 175: 1417-1422.
193. Kullberg, M. C., E. J. Pearce, S. E. Hieny, A. Sher, and J. A. Berzofsky. 1992. Infection with *Schistosoma mansoni* alters Th1/Th2 cytokine responses to a non-parasite antigen. *J. Immunol.* 148: 3264-3270.
194. Sher, A., R. T. Gazzinelli, I. P. Oswald, M. Clerici, M. Kullberg, E. J. Pearce, J. A. Berzofsky, T. R. Mosmann, S. L. James, H. C. Morse, III, and G. M. Shearer. 1992. Role of T-cell derived cytokines in the downregulation of immune responses in parasitic and retroviral infection. *Immunological Reviews* 127: 183-204.
195. Shirai, M., T. Akatsuka, C. D. Pendleton, R. Houghten, C. Wychowski, K. Mihalik, S. Feinstone, and J. A. Berzofsky. 1992. Induction of cytotoxic T cells to a crossreactive epitope in the hepatitis C viral nonstructural RNA polymerase-like protein. *J. Virol.* 66: 4098-4106.
196. Clerici, M., J. A. Berzofsky, G. M. Shearer, J. V. Giorgi, and C. Tacket. 1992. On HIV-serologic testing of blood and tissue donors. *New Engl. J. Med.* 327: 564-565 (letter).

197. Quakyi, I. A., D. W. Taylor, A. H. Johnson, J. B. Allotey, J. A. Berzofsky, L. H. Miller, and M. F. Good. 1992. Development of a malaria T-cell vaccine for blood stage immunity. *Scand. J. Immunol.* 36 (Suppl. 11):9-16.
198. Hosmalin, A., S. Kumar, D. Barnd, R. Houghten, G. E. Smith, S. H. Hughes, and J. A. Berzofsky. 1992. Immunization with soluble protein-pulsed spleen cells induces class I-restricted CTL that recognize immunodominant epitopic peptides from *P. falciparum* and HIV-1. *J. Immunol.* 149: 1311-1318.
199. Weiss, W. R., J. A. Berzofsky, R. A. Houghten, M. Sedegah, M. Hollindale, and S. L. Hoffman. 1992. A T cell clone directed at the circumsporozoite protein which protects mice against both *Plasmodium yoelii* and *Plasmodium berghei*. *J. Immunol.* 149: 2103-2109.
200. Actor, J. K., M. Shirai, M. C. Kullberg, R. M. L. Buller, A. Sher, and J. A. Berzofsky. 1993. Helminth infection results in decreased virus-specific CD8⁺ cytotoxic T-cell and Th1 cytokine responses as well as delayed virus clearance. *Proc. Natl. Acad. Sci. USA* 90: 948-952.

201. Boehncke, W. -H., T. Takeshita, C. D. Pendleton, S. Sadegh-Nasseri, L. Racioppi, R. A. Houghten, J. A. Berzofsky, and R. N. Germain. 1993. The importance of dominant negative effects of amino acids side chain substitution in peptide-MHC molecule interactions and T cell recognition. *J. Immunol.* 150: 331-341.
202. Takahashi, H., Y. Nakagawa, M. Takeuchi, K. Yokomuro, and J. A. Berzofsky. 1993. Elicitation of CD8⁺ class-I-restricted CTLs by immunization with syngeneic irradiated HIV-1 envelope-derived peptide-pulsed splenic dendritic cells. In *Vaccines 93*. F. Brown, R.M. Chanock, H.S. Ginsberg, and R.A. Lerner, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
203. Berzofsky, J. A. and I. J. Berkower. 1993. Immunogenicity and antigen structure. In *Fundamental Immunology*. 3rd Edn. W.E. Paul, editor. Raven Press, New York., pp. 235-282.
204. Berzofsky, J. A., I. J. Berkower, and S. L. Epstein. 1993. Antigen-antibody interactions and monoclonal antibodies. In *Fundamental Immunology*. 3rd Edn. W.E. Paul, editor. Raven Press, New York. , pp. 421-465.
205. Berzofsky, J. A. 1995. Immunogenicity and Antigenicity. In *Samter's Immunological Diseases* . M.M. Frank, K.F. Austen, H.N. Claman, and E.R. Unanue, editors. Little, Brown, and Company, Boston, pp. 17-32..
206. Berzofsky, J. A. 1993. Epitope selection and design of synthetic vaccines: molecular approaches to enhancing immunogenicity and crossreactivity of engineered vaccines. *Annals of the New York Academy of Sciences* , 690: 256-264.
207. Ahlers, J. D., C. D. Pendleton, N. Dunlop, A. Minassian, P. L. Nara, and J. A. Berzofsky. 1993. Construction of an HIV-1 peptide vaccine containing a multideterminant helper peptide linked to a V3 loop peptide 18 inducing strong neutralizing antibody responses in mice of multiple MHC haplotypes following two immunizations. *J. Immunol.* 150: 5647-5665.
208. Wasserman, G. M., S. Kumar, J. Ahlers, F. Ramsdell, J. A. Berzofsky, and L. H. Miller. 1993. An approach to development of specific T lymphocyte lines using preprocessed antigens in murine malaria *Plasmodium vinckei vinckei*. *Infect. Immun.* 61: 1958-1963.
209. Takahashi, H., Y. Nakagawa, K. Yokomuro, and J. A. Berzofsky. 1993. Induction of CD8⁺ CTL by immunization with syngeneic irradiated HIV-1 envelope derived peptide-pulsed dendritic cells. *Internat. Immunol.* 5: 849-857.
210. Shirai, M., M. S. Vacchio, R. J. Hodes, and J. A. Berzofsky. 1993. Preferential V β usage by cytotoxic T cells crossreactive between two epitopes of HIV-1 gp160 and degenerate in class I MHC restriction. *J. Immunol.* 151: 2283-2295.

211. Yanuck, M., D. P. Carbone, C. D. Pendleton, T. Tsukui, S. F. Winter, J. D. Minna, and J. A. Berzofsky. 1993. A mutant p53 tumor suppressor protein is a target for peptide-induced CD8⁺ cytotoxic T cells. *Cancer. Res.* 53: 3257-3261.

212. Takeshita, T., S. Kozlowski, R. D. England, R. Brower, J. Schneck, H. Takahashi, C. DeLisi, D. H. Margulies, and J. A. Berzofsky. 1993. Role of conserved regions of class I MHC molecules in the activation of CD8⁺ CTL by peptide and purified cell-free class I molecules. *Internat. Immunol.* 5: 1129-1138.
213. Kozlowski, S., M. Corr, M. Shirai, L. F. Boyd, C. D. Pendleton, J. A. Berzofsky, and D. H. Margulies. 1993. Multiple pathways are involved in the extracellular processing of major histocompatibility complex class I-restricted peptides. *J. Immunol.* 151: 4033-4044.
214. Williams, M. E., M. C. Kullberg, S. Barbieri, P. Caspar, J. A. Berzofsky, R. A. Seder, and A. Sher. 1993. FcεR⁺ cells are a major source of antigen induced IL-4 in spleens of mice infected with *Schistosoma mansoni*. *Eur. J. Immunol.* 23: 1910-1916.
215. Shirai, M., C. D. Pendleton, J. Ahlers, T. Takeshita, M. Newman, and J. A. Berzofsky. 1994. Helper-CTL determinant linkage required for priming of anti-HIV CD8⁺ CTL in vivo with peptide vaccine constructs. *J. Immunol.* 152: 549-556.
216. Clerici, M., A. V. Sison, J. A. Berzofsky, T. A. Rakusan, C. D. Brandt, M. Ellaurie, M. Villa, C. Colie, D. J. Venzon, J. L. Sever, and G. M. Shearer. 1993. Cellular immune factors associated with mother-to-infant transmission of HIV. *AIDS* 7:1427-1433.
217. Clerici, M., D. R. Lucey, J. A. Berzofsky, L. A. Pinto, T. A. Wynn, S. P. Blatt, M. J. Dolan, C. W. Hendrix, S. F. Wolf, and G. M. Shearer. 1993. Restoration of HIV-specific cell-mediated immune responses by interleukin-12 *in vitro*. *Science* 262: 1721-1724.
218. Clerici, M., T. A. Wynn, J. A. Berzofsky, S. P. Blatt, C. W. Hendrix, A. Sher, R. L. Coffman, and G. M. Shearer. 1994. Role of interleukin-10 (IL-10) in T helper cell dysfunction in asymptomatic individuals infected with the human immunodeficiency virus (HIV-1). *J. Clin. Invest.* 93: 768-775.
219. Cease, K. B. and J. A. Berzofsky. 1994. Towards a vaccine for AIDS: the emergence of immunobiology-based vaccine development. *Annu. Rev. Immunol.* 12: 923-989.
220. Altuvia, Y., J. A. Berzofsky, R. Rosenfeld, and H. Margalit. 1994. Sequence features that correlate with MHC restriction. *Molec. Immunol.* 31: 1-19.
221. Berzofsky, J. A., J. D. Ahlers, M. Shirai, C. D. Pendleton, T. Takeshita, N. Dunlop, A. Minassian, M. Newman, and P. L. Nara. 1994. Construction of candidate synthetic peptide vaccines for HIV-1. In *Vaccines 94: Modern approaches to new vaccines including prevention of AIDS*. E. Norrby, F. Brown, R.M. Chanock, and H.S. Ginsberg, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. pp. 147-153.

222. Takahashi, H., T. Takeshita, S. Kozlowski, Y. Nakagawa, J. Ahlers, C. D. Pendleton, R. L. Moore, B. S. Fox, K. Yokomuro, D. H. Margulies, and J. A. Berzofsky. 1994. An immunodominant determinant of HIV-1 envelope recognized by both class-I-restricted CD8⁺ CTLs and class II-restricted CD4⁺ helper T cells shares similar MHC-binding sites. In *Vaccines 94: Modern approaches to new vaccines including prevention of AIDS*. E. Norrby, F. Brown, R.M. Chanock, and H.S. Ginsberg, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor. pp. 181-188.
223. Clerici, M., J. M. Levin, H. A. Kessler, A. Harris, J. A. Berzofsky, A. L. Landay, and G. M. Shearer. 1994. HIV-specific T-helper activity in seronegative health care workers exposed to contaminated blood. *J. Amer. Med. Assoc.* 271:42-46.
224. Clerici, M. and J. A. Berzofsky. 1994. Cellular immunity and cytokines in HIV infection. *AIDS* 8: S175-S182.
225. Wiedenfeld, E. A., M. Fernandez-Viña, J. A. Berzofsky, and D. P. Carbone. 1994. Evidence for selection against human lung cancers bearing p53 missense mutations which occur within the HLA A *0201 peptide consensus motif. *Cancer Res.* 54:1175-1177.
226. Weissler, J. C., L. E. Rosenberg, and J. A. Berzofsky. 1994. The spring meetings: Back to the future. *J. Clin. Invest.* 93:457-458.
227. Shirai, M., H. Okada, M. Nishioka, T. Akatsuka, C. Wychowski, R. Houghten, C. D. Pendleton, S. M. Feinstone, and J. A. Berzofsky. 1994. An epitope in hepatitis C virus core region recognized by cytotoxic T cells in mice and humans. *J. Virol.* 68:3334-3342.
228. Kozhich, A. T., Y. -I. Kawano, C. E. Egwuagu, R. R. Caspi, R. K. Maturi, J. A. Berzofsky, and I. Gery. 1994. A pathogenic autoimmune process targeted at a surrogate epitope. *J. Exp. Med.* 180: 133-140.
229. Bermas, B. L., M. Petri, J. A. Berzofsky, A. Waisman, G. M. Shearer, and E. Mozes. 1994. Autoantibodies in mice with experimentally induced systemic lupus erythematosus (SLE) and patients with SLE bind gp120 and peptides from the HIV-1 envelope. *AIDS Res. Hum. Retroviruses.* 10: 1071-1077.
230. Berzofsky, J. A. 1994. Presidential address to the American Society for Clinical Investigation, Baltimore, Maryland, 30 April, 1994: Cross-fertilization among fields: a seminal event in the progress of biomedical research. *J. Clin. Invest.* 94: 911-918.
231. Brower, R. C., R. England, T. Takeshita, S. Kozlowski, D. H. Margulies, J. A. Berzofsky, and C. DeLisi. 1994. Minimal requirements for peptide mediated activation of CD8⁺ CTL. *Molec. Immunol.* 31: 1285-1293.
232. Quakyi, I. A., J. Currier, A. Fell, D. W. Taylor, T. Roberts, R. A. Houghten, R. D. England, J. A. Berzofsky, L. H. Miller, and M. F. Good. 1994. Analysis of human

T cell clones specific for conserved peptide sequences within malaria proteins: paucity of clones responsive to intact parasites. *J. Immunol.* 153: 2082-2092.

233. Nishimura, M., A. G. Kermode, M. Clerici, G. M. Shearer, J. A. Berzofsky, T. Uchiyama, S. Z. Wiktor, E. Pate, B. Maloney, A. Manns, W. Blattner, and S. Jacobson. 1994. Demonstration of human T lymphotropic virus type (HTLV-I)-specific T cell responses from seronegative and polymerase chain reaction-negative persons exposed to HTLV-I. *J. Infect. Dis.* 170:334-338.
234. Actor, J. K., M. A. Marshall, I. A. Eltoun, R. M. L. Buller, J. A. Berzofsky, and A. Sher. 1994. Increased susceptibility of mice infected with *Schistosoma mansoni* to recombinant vaccinia virus: association of viral persistence with egg granuloma formation. *Eur. J. Immunol.* 24: 3050-3056.
235. Meister, G. E., C. G. P. Roberts, J. A. Berzofsky, and A. S. DeGroot. 1995. Two novel T cell epitope prediction algorithms based on MHC-binding motifs; comparison of predicted and published epitopes from *Mycobacterium tuberculosis* and HIV protein sequences. *Vaccine* , 13: 581-591.
236. Takeshita, T., H. Takahashi, S. Kozlowski, J. D. Ahlers, C. D. Pendleton, R. L. Moore, Y. Nakagawa, K. Yokomuro, B. S. Fox, D. H. Margulies, and J. A. Berzofsky. 1995. Molecular analysis of the same HIV peptide functionally binding to both a class I and a class II MHC molecule. *J. Immunol.* 154: 1973-1986.
237. Berzofsky, J. A., J. D. Ahlers, M. Alexander-Miller, T. Tsukui, C. D. Pendleton, N. Dunlop, D. P. Carbone, and P. L. Nara. 1995. Developing synthetic peptide vaccines for HIV-1. In *Molecular approaches to the control of infectious diseases*. F. Brown, R. Chanock, H.S. Ginsberg, and E. Norrby, editors. Cold Spring Harbor Laboratory, Cold Spring Harbor, 135-142.
238. Berzofsky, J. A. and I. J. Berkower. 1995. Novel approaches to peptide and engineered protein vaccines for HIV using defined epitopes: advances in 1994-95. *AIDS* 9: S143-157.
239. Shirai, M., T. Arichi, M. Nishioka, T. Nomura, K. Ikeda, K. Kawanishi, V. H. Engelhard, S. M. Feinstone, and J. A. Berzofsky. 1995. Cytotoxic T lymphocyte (CTL) responses of HLA-A2.1-transgenic mice specific for hepatitis C viral peptides predict epitopes for CTL of humans carrying HLA-A2.1. *J. Immunol.* 154: 2733-2742.
240. Battegay, M., J. Fikes, A. M. Di Bisceglie, P. A. Wentworth, A. Sette, E. Celis, W. -M. Ching, A. Grakoui, C. M. Rice, K. Kurokohchi, J. A. Berzofsky, J. H. Hoofnagle, S. M. Feinstone, and T. Akatsuka. 1995. Patients with chronic hepatitis C have circulating cytotoxic T cells which recognize hepatitis C virus-encoded peptides binding to HLA-A2.1 molecules. *J. Virol.* 69:2462-2470.
241. Berzofsky, J. A. 1995. Designing peptide vaccines to broaden recognition and enhance potency. *Ann. N. Y. Acad. Sci.* 754: 161-168.

242. Malik, A., R. Houghten, G. Corradin, S. Buus, J. A. Berzofsky, and S. L. Hoffman.
1995. Identification of a nonameric H-2K^k-restricted CD8⁺ cytotoxic lymphocyte
epitope on the *Plasmodium falciparum* circumsporozoite protein. *Infect. Immun.* 63:
1955-1959.

243. Pinto, L. A., J. Sullivan, J. A. Berzofsky, M. Clerici, H. A. Kessler, A. L. Landay, and G. M. Shearer. 1995. *Env*-specific cytotoxic T lymphocyte responses in HIV seronegative health care workers occupationally exposed to HIV-contaminated body fluids. *J. Clin. Invest.* 96: 867-876.
244. Cornette, J. L., H. Margalit, J. A. Berzofsky, and C. DeLisi. 1995. Periodic variation in side chain polarities of T cell antigenic peptides correlates with their structure and activity. *Proc. Nat. Acad. Sci. USA.* 92: 8368-8372.
245. England, R. E., M. C. Kullberg, J. L. Cornette, and J. A. Berzofsky. 1995. Molecular analysis of a heteroclitic T-cell response to the immunodominant epitope of sperm whale myoglobin: implications for peptide partial agonists. *J. Immunol.* 155: 4295-4306.
246. Shirai, M., M. Chen, T. Arichi, M. Nishioka, M. Newman, T. Nakazawa, S. M. Feinstone, and J. A. Berzofsky. 1996. Use of intrinsic and extrinsic helper epitopes for in vivo induction of anti-hepatitis C virus cytotoxic T lymphocytes (CTL) with CTL epitope peptide vaccines. *J. Infect. Dis.* 173: 24-31.
247. Shearer, G. M., M. Clerici, A. Sarin, J. A. Berzofsky, and P. A. Henkart. 1995. Cytokines in immune regulation/pathogenesis in HIV infection. Ciba Foundation Symposium 195: 142-153 (Wiley, Chichester).
248. Seder, R. A., K. H. Grabstein, J. A. Berzofsky, and J. F. McDyer. 1995. Cytokine interactions in human immunodeficiency virus-infected individuals: Roles of interleukin (IL)-2, IL-12, and IL-15. *J. Exp. Med.* 182:1067-1078.
249. Kurokohchi, K., T. Akatsuka, C. D. Pendleton, A. Takamizawa, M. Nishioka, M. Battegay, S. M. Feinstone, and J. A. Berzofsky. 1996. Use of recombinant protein to identify a motif-negative human CTL epitope presented by HLA-A2 in the hepatitis C virus NS3 region. *J. Virol.* 70: 232-240.
250. Takahashi, H., Y. Nakagawa, G. Leggatt, Y. Ishida, T. Saito, K. Yokomuro, and J. A. Berzofsky. 1996. Inactivation of HIV-1 envelope-specific CD8⁺ cytotoxic T lymphocytes by free antigenic peptide: A self-veto mechanism? *J. Exp. Med.* 183: 879-889.
251. Ahlers, J. D., N. Dunlop, C. D. Pendleton, M. Newman, P. L. Nara, and J. A. Berzofsky. 1996. Candidate HIV-1 multideterminant cluster peptide-P18MN vaccine constructs elicit Th1 helper T cells, cytotoxic T cells, and neutralizing antibody, all using the same adjuvant immunization. *AIDS Res. Hum. Retroviruses* 12: 259-272.
252. Kurokohchi, K., M. Carrington, D. L. Mann, T. B. Simonis, S. M. Feinstone, T. Akatsuka, and J. A. Berzofsky. 1996. Expression of HLA class I molecules and the transporter associated with antigen precessing (TAP) in hepatocellular carcinoma. *Hepatology* 23: 1181-1188.

253. Kullberg, M. C., J. A. Berzofsky, D. Lj. Jankovic, S. Barbieri, M. E. Williams, P. Perlmann, A. Sher, and M. Troye-Blomberg. 1996. T-cell-derived IL-3 induces the production of IL-4 by non-B, non-T cells to amplify the Th2-cytokine response to a non-parasite antigen in *Schistosoma mansoni*-infected mice. *J. Immunol.* 156: 1482-1489.
254. Quakyi, I. A., L. H. Miller, M. F. Good, J. D. Ahlers, S. N. Isaacs, J. H. Nunberg, R. A. Houghten, D. B. Keister, J. E. Coligan, B. Moss, D. Alling, J. A. Berzofsky, and D. C. Kaslow. 1995. Synthetic Peptides from *P. falciparum* sexual stage 25-kDa protein induce antibodies that react with the native protein: the role of IL-2 and conformational structure on immunogenicity of Pfs25. *Peptide Research* 8: 335-344.
255. Ciernik, I. F., J. A. Berzofsky, and D. P. Carbone. 1995. Mutant oncoprotein immunization induces CTL specifically lysing tumor cells endogenously expressing the corresponding intact mutant p53. *Hybridoma* 14:139-142.
256. Alexander-Miller, M. A., K. C. Parker, T. Tsukui, C. D. Pendleton, J. E. Coligan, and J. A. Berzofsky. 1996. Molecular analysis of presentation by HLA-A2.1 of a promiscuously binding V3 loop peptide from the HIV-1 envelope protein to human CTL. *Int. Immunol.* 8: 641-649.
257. Alexander-Miller, M. A., G. R. Leggatt, and J. A. Berzofsky. 1996. Selective expansion of high or low avidity cytotoxic T lymphocytes and efficacy for adoptive immunotherapy. *Proc. Natl. Acad. Sci. U. S. A.* 93: 4102-4107.
258. Roberts, C. G. P., G. E. Meister, B. T. Jesdale, J. Lieberman, J. A. Berzofsky, and A. S. DeGroot. 1996. Prediction of HIV peptide epitopes by a novel algorithm. *AIDS Res. Hum. Retroviruses* 12: 593-610.
259. Landay, A. L., M. Clerici, F. Hashemi, H. Kessler, J. A. Berzofsky, and G. M. Shearer. 1996. In vitro restoration of T cell immune function in human immunodeficiency virus-positive persons: effects of interleukin (IL)-12 and anti-IL-10. *J. Infect. Dis.* 173: 1085-1091.
260. Ciernik, I. F., J. A. Berzofsky, and D. P. Carbone. 1996. Human lung cancer cells endogenously expressing mutant p53 process and present the mutant epitope, and are lysed by mutant-specific CTL. *Clin. Cancer. Res.* 2: 877-882..
261. Ciernik, I. F., J. A. Berzofsky, and D. P. Carbone. 1996. Induction of cytotoxic T lymphocytes and anti-tumor immunity with DNA vaccines expressing single T cell epitopes. *J. Immunol.* 156: 2369-2375.
262. Gabrilovich, D. I., S. Nadaf, J. Corak, J. A. Berzofsky, and D. P. Carbone. 1996. Dendritic cells in anti-tumor immune responses. II. Dendritic cells grown from bone marrow precursors, but not mature DC from tumor-bearing mice are effective antigen carriers in the therapy of established tumors.. *Cellular Immunol.* 170: 111-119.

263. Clerici, M., A. Sarin, J. A. Berzofsky, A. L. Landay, H. A. Kessler, F. Hashemi, C. W. Hendrix, S. P. Blatt, J. Rusnak, M. J. Dolan, R. L. Coffman, P. A. Henkart, and G. M. Shearer. 1996. Antigen-stimulated apoptotic T cell death in HIV infection is selective for CD4⁺ T cells, modulated by cytokines and effected by lymphotoxin. *AIDS* 10: 603-611.
264. Tian, J. -H., L. H. Miller, D. C. Kaslow, J. Ahlers, M. F. Good, D. W. Alling, J. A. Berzofsky, and S. Kumar. 1996. Genetic regulation of protective immune response in congenic strains of mice vaccinated with a subunit malaria vaccine. *J. Immunol.* 157: 1176-1183.
265. Alexander-Miller, M. A., G. R. Leggatt, A. Sarin, and J. A. Berzofsky. 1996. Role of antigen, CD8, and CTL avidity in high dose antigen induction of apoptosis of effector CTL. *J. Exp. Med.* 184: 485-492.
266. Tsukui, T., A. Hildesheim, M. H. Schiffman, J. Lucci III, D. Contois, P. Lawler, B. B. Rush, A. T. Lorincz, A. Corrigan, R. D. Burk, W. Qu, M. A. Marshall, D. Mann, M. Carrington, M. Clerici, G. M. Shearer, D. P. Carbone, D. R. Scott, R. A. Houghten, and J. A. Berzofsky. 1996. IL-2 production in vitro by peripheral lymphocytes in response to human papillomavirus-derived peptides: correlation with cervical pathology. *Cancer Res.* 56: 3967-3974.
267. Huang, X., M. C. Smith, J. A. Berzofsky, and J. J. Jr. Barchi. 1996. Structural comparison of a 15 residue peptide from the V3 loop of HIV-1_{IIIB} and an O-glycosylated analogue. *FEBS Letters* 393: 280-286.
268. Shirai, M., K. Kurokohchi, C. D. Pendleton, T. Arichi, L. F. Boyd, H. Takahashi, D. H. Margulies, and J. A. Berzofsky. 1996. Reciprocal CTL cross-reactivity interactions between two major epitopes within HIV-1 gp160. *J. Immunol.* , 157: 4399-4411.
269. Leggatt, G. R., M. A. Alexander-Miller, A. Kumar (née Malik), S. L. Hoffman, and J. A. Berzofsky. 1997. Cytotoxic T lymphocyte (CTL) adherence assay (CAA): A non-radioactive assay for murine CTL recognition of peptide-MHC class I complexes. *J. Immunol. Methods*, 201: 1-10.
270. Sarin, A., M. S. Williams, M. A. Alexander-Miller, J. A. Berzofsky, C. M. Zacharchuk, and P. A. Henkart. 1997. Target cell lysis by CTL granule exocytosis is independent of ICE/Ced-3 family proteases. *Immunity* 6: 209-215.
271. Saito, T., G. J. Sherman, K. Kurokohchi, Z. -P. Guo, M. Donets, M. -Y. W. Yu, J. A. Berzofsky, T. Akatsuka, and S. M. Feinstone. 1997. Plasmid DNA-based immunization for hepatitis C virus structural proteins, immune responses in mice. *Gastroenterology* 112: 1321-1330.
272. Shirai, M., S. Kozlowski, D. H. Margulies, and J. A. Berzofsky. 1997. Degenerate MHC restriction reveals the contribution of class I MHC molecules in determining the fine

- specificity of CTL recognition of an immunodominant determinant of HIV-1 gp160 V3 loop. *J. Immunol.* 158: 3181-3188.
273. Ahlers, J. D., N. Dunlop, D. W. Alling, P. L. Nara, and J. A. Berzofsky. 1997. Cytokine-in-adjuvant steering of the immune response phenotype to HIV-1 vaccine constructs: GM-CSF and TNF α synergize with IL-12 to enhance induction of CTL. *J. Immunol.* 158: 3947-3958.
 274. Kozhich, A. T., R. R. Caspi, J. A. Berzofsky, and I. Gery. 1997. Immunogenicity and immunopathogenicity of an autoimmune epitope are potentiated by increasing MHC binding through residue substitution. *J. Immunol.* 158: 4145-4151.
 275. Zhang, C., J. L. Cornette, J. A. Berzofsky, and C. DeLisi. 1997. The organization of human leukocyte antigen class I epitopes in HIV genome products: Implications for HIV evolution and vaccine design. *Vaccine* 15: 1291-1302.
 276. Smith, M. C., C. D. Pendleton, V. E. Maher, M. J. Kelley, D. P. Carbone, and J. A. Berzofsky. 1997. Oncogenic mutations in ras create HLA-A2.1 binding peptides but affect their extracellular processing. *Internat. Immunol.* 9: 1085-1093.
 277. Goletz, T. J., K. R. Klimpel, S. H. Leppla, J. M. Keith, and J. A. Berzofsky. 1997. Delivery of antigens to the MHC class I pathway using bacterial toxins. *Hum. Immunol.* 54: 129-136..
 278. DeGroot, A. S., B. M. Jesdale, and J. A. Berzofsky. 1998. Prediction and determination of MHC ligands and T cell epitopes. In *Immunological Methods in Microbiology* Vol. 25. Immunology of Infection. S.H.E. Kaufmann and D. Kabelitz, editors. Academic Press, London, Chap. 3, pp. 79-106.
 279. Hirunpetcharat, C., J. -H. Tian, D. C. Kaslow, N. van Rooijen, S. Kumar, J. A. Berzofsky, L. H. Miller, and M. F. Good. 1997. Complete protective immunity induced in mice by immunization with the 19kDa carboxyterminal fragment of the merozoite surface protein-1 (MSP1₁₉) of *Plasmodium yoelii* expressed in *Saccharomyces cerevisiae*: Correlation of protection with antigen-specific antibody titer, but not effector CD4⁺ T cells. *J. Immunol.* 159: 3400-3411.
 280. DeGroot, A. S., G. E. Meister, J. L. Cornette, H. Margalit, C. DeLisi, and J. A. Berzofsky. 1997. Computer prediction of T-cell epitopes. In *New Generation Vaccines*. M.M. Levine, G.C. Woodrow, J.B. Kaper, and G.S. Cobon, editors. Marcel Dekker, Inc., New York. 127-138.
 281. Ahlers, J. D., T. Takeshita, C. D. Pendleton, and J. A. Berzofsky. 1997. Enhanced immunogenicity of HIV-1 vaccine construct by modification of the native peptide sequence. *Proc. Natl. Acad. Sci. U. S. A.* 94: 10856-10861.

282. Shirai, M., T. Arichi, T. Nakazawa, and J. A. Berzofsky. 1998. Persistent infection by Helicobacter pylori downmodulates virus-specific CD8⁺ cytotoxic T cell response and prolongs viral infection. *J. Infect. Dis.* 177: 72-80.
283. Goletz, T.J., K. R. Klimpel, N. Arora, S. H. Leppla, J. M. Keith, and J. A. Berzofsky. 1997. Targeting HIV proteins to the major histocompatibility complex class I processing pathway with a novel gp120-anthrax toxin fusion protein. *Proc. Natl. Acad. Sci. U.S.A.* 94: 12059-12064.
284. Belyakov, I. M., M. A. Derby, J. D. Ahlers, B. L. Kelsall, P. Earl, B. Moss, W. Strober, and J. A. Berzofsky. 1998. Mucosal immunization with HIV-1 peptide vaccine induces mucosal and systemic cytotoxic T lymphocytes and protective immunity in mice against intrarectal recombinant HIV-vaccinia challenge. *Proc. Natl. Acad. Sci.* 95:1709-1714.
285. Hildesheim, A., M. H. Schiffman, T. Tsukui, C. A. Swanson, J. Lucci, D. R. Scott, A. G. Glass, B. B. Rush, A. T. Lorincz, A. Corrigan, R. D. Burk, K. Helgesen, R. A. Houghten, M. E. Sherman, R. J. Kurman, J. A. Berzofsky, and T. R. Kramer. 1997. Immune activation in cervical neoplasia: Cross-sectional association between plasma soluble interleukin 2 receptor levels and disease. *Cancer Epidemiol. Biomark. Preven.* 6:807-813.
286. Leggatt, G. R. and J. A. Berzofsky. 1999. Cytotoxic T cell adherence assay (CAA). In *Methods in Molecular Biology Series : T cell protocols*. K. Kearse, editor. Humana Press, Totowa, NJ., 134: 277-281.

287. Belyakov, I. M., L. S. Wyatt, J. D. Ahlers, P. Earl, C. D. Pendleton, B. L. Kelsall, W. Strober, B. Moss, and J. A. Berzofsky. 1998. Induction of mucosal CTL response by intrarectal immunization with a replication-deficient recombinant vaccinia virus expressing HIV 89.6 envelope protein. *J. Virol.* 72: 8264-8272.
288. Leggatt, G. R., A. Hosmalin, C. D. Pendleton, A. Kumar, S. Hoffman, and J. A. Berzofsky. 1998. The importance of pairwise interactions between peptide residues in the delineation of T cell receptor specificity. *J Immunol.* 161: 4728-4735:
289. Sarobe, P., C. D. Pendleton, T. Akatsuka, D. Lau, V. H. Engelhard, S. M. Feinstone, and J. A. Berzofsky. 1998. Enhanced in vitro potency and in vivo immunogenicity of a CTL epitope from hepatitis C virus core protein following amino acid replacement at secondary HLA-A2.1 binding positions. *J. Clin. Invest.* 102: 1239-1248.
290. Alexander-Miller, M. A., M. A. Derby, A. Sarin, P. A. Henkart, and J. A. Berzofsky. 1998. Supra-optimal peptide/MHC causes a decrease in Bcl-2 and allows TNF- α receptor II-mediated apoptosis of CTL. *J. Exp. Med.* 188: 1391-1399.
291. Berzofsky, J. A., I. J. Berkower, and S. L. Epstein. 1999. Antigen-antibody interactions and monoclonal antibodies. In *Fundamental Immunology*. W.E. Paul, editor. Lippincott-Raven, Philadelphia. 75-110.
292. Berzofsky, J. A. and I. J. Berkower. 1999. Immunogenicity and antigen structure. In *Fundamental Immunology*. W.E. Paul, editor. Lippincott-Raven, Philadelphia. 651-699.
293. Goletz, T. J., C. L. Mackall, J. A. Berzofsky, and L. J. Helman. 1998. Molecular alterations in pediatric sarcomas: potential targets for immunotherapy. *Sarcoma* 2:77-87.
294. Shirai, M., T. Arichi, M. Chen, M. Nishioka, K. Ikeda, H. Takahashi, N. Enomoto, T. Saito, M. E. Major, T. Nakazawa, T. Akatsuka, S. M. Feinstone, and J. A. Berzofsky. 1999. T cell recognition of hypervariable region 1 from hepatitis C virus envelope protein with multiple class II MHC molecules in mice and humans: Preferential help for induction of antibodies to the hypervariable region.. *J. Immunol.* 162: 568-576.
295. Oscherwitz, J., F. M. Gotch, K. B. Cease, and J. A. Berzofsky. 1999. New insights and approaches regarding B and T cell epitopes in HIV vaccine design. *AIDS* , 13 (Suppl. A): S163-S174.
296. Suh, W-K., M. A. Derby, M. F. Cohen-Doyle, G. J. Schoenhals, K. Fröh, J. A. Berzofsky, and D. B. Williams. 1999. Interaction of murine MHC class I molecules with tapasin and TAP enhances peptide loading and involves the heavy chain $\alpha 3$ domain. *J. Immunol.* 162: 1530-1540.

297. Belyakov, I. M., J. D. Ahlers, B. Y. Brandwein, P. Earl, B. L. Kelsall, B. Moss, W. Strober, and J. A. Berzofsky. 1998. The importance of local mucosal HIV-specific CD8⁺ cytotoxic T lymphocytes for resistance to mucosal-viral transmission in mice and enhancement of resistance by local administration of IL-12. *J. Clin. Invest.* 102: 2072-2081.
298. D'Amico, R., L. A. Pinto, P. Meyer, A. L. Landay, A. A. Harris, M. Clerici, J. A. Berzofsky, G. M. Shearer, and H. A. Kessler. 1999. Effect of zidovudine postexposure prophylaxis on the development of HIV-specific cytotoxic T-lymphocyte responses in HIV-exposed health care workers. *Infection Control and Hospital Epidemiology*, 20: 428-430.
299. McDyer, J. F., M. Dybul, T. J. Goletz, A. L. Kinter, E. K. Thomas, J. A. Berzofsky, A. S. Fauci, and R. A. Seder. 1999. Differential effects of CD40 ligand/trimer stimulation on the ability of dendritic cells to replicate and transmit HIV infection: Evidence for CC-chemokine-dependent and -independent mechanisms. *J. Immunol.*, 162: 3711-3717.
300. Belyakov, I. M., B. Moss, W. Strober, and J. A. Berzofsky. 1999. Mucosal vaccination overcomes the barrier to recombinant vaccinia immunization caused by preexisting poxvirus immunity. *Proc. Natl. Acad. Sci. U. S. A.* 96: 4512-4517.
301. Berzofsky, J. A., J. D. Ahlers, M. A. Derby, C. D. Pendleton, T. Arichi, and I. M. Belyakov. 1999. Approaches to improve engineered vaccines for HIV and other viruses that cause chronic infections. *Immunological Reviews* 170: 151-172.
302. Matsui, S., J. D. Ahlers, A. O. Vortmeyer, M. Terabe, T. Tsukui, D. P. Carbone, L. A. Liotta, and J. Berzofsky. 1999. A model for CD8⁺ CTL tumor immunosurveillance and regulation of tumor escape by CD4⁺ T cells through an effect on quality of CTL. *J. Immunol.* 163: 184-193.
303. Maher, V. E., B. S. Worley, D. Contois, M. C. Smith, M. J. Kelley, M. Stipanov, S. N. Khleif, T. Goletz, L. van den Broeke, C. Mackall, L. J. Helman, D. P. Carbone, and J. A. Berzofsky. 2000. Mutant oncogene and tumor suppressor gene products and fusion proteins created by chromosomal translocations as targets for cancer vaccines. In *Peptide-based cancer vaccines*. W.M. Kast, editor. Landes Bioscience, Austin, pp. 17-39.
304. Chung, D. H., J. Dorfman, D. Plaksin, K. Natarajan, I. M. Belyakov, R. Hunziker, J. A. Berzofsky, W. M. Yokoyama, M. G. Mage, and D. H. Margulies. 1999. Natural killer (NK) and cytolytic T lymphocyte (CTL) recognition of a single chain H-2D^d molecule: distinct sites of H-2D^d interact with NK and T cell receptors. *J. Immunol.* 163: 3699-3708.
305. Pinto, L. A., J. A. Berzofsky, K. R. Fowke, R. F. Littles, F. Merced-Galindez, R. Humphreys, J. Ahlers, N. Dunlop, R. B. Cohen, S. M. Steinberg, P. Nara, G. M. Shearer, and R. Yarchoan. 1999. HIV-specific immunity following immunization

with HIV synthetic envelope peptides in asymptomatic HIV-infected patients. *AIDS* 13: 2003-2012.

306. Berzofsky, J. A., L. J. Helman, and D. P. Carbone. 2001. Oncogene products and mutated proteins as tumor antigens. In *Biologic Therapy of Cancer: Principles and Practice* (3rd Ed.). S. A. Rosenberg, editor. Lippincott Williams & Wilkins, Philadelphia, in press.
307. Arichi, T., T. Saito, M. E. Major, I. M. Belyakov, M. Shirai, V. H. Engelhard, S. M. Feinstone, and J. A. Berzofsky. 2000. Prophylactic DNA vaccine for hepatitis C virus (HCV) infection: HCV specific CTL induction and protection from HCV-recombinant vaccinia infection in an HLA-A2.1 transgenic mouse model. *Proc. Natl. Acad. Sci. U.S.A.* 97: 297-302. (Retracted due to discovery of fraud by first author.)
308. Ciernik, I. F., P. Romero, J. A. Berzofsky, and D. P. Carbone. 1999. Ionizing radiation enhances immunogenicity of cells expressing a tumor-specific T-cell epitope. *Int. J. Rad. Oncolo. Biol. Phys.* 45:735-741.
309. Parra, M., G. Hui, A. Johnson, J. A. Berzofsky, T. Roberts, I. A. Quakyi, and D. W. Taylor. 2000. Characterization of conserved T and B cell epitopes in *Plasmodium falciparum* major merozoite surface protein-1 (MSP-1). *Infect. Immun.* 68: 2685-2691.
310. Berzofsky, J. A. 2000. Immunogenicity and Antigenicity. In *Samter's Immunologic Diseases* Sixth Edition. K.F. Austen, M.M. Frank, J.P. Atkinson, and H. Cantor, editors. Lippincott Williams & Wilkins, Baltimore.
311. Berzofsky J.A., J.D. Ahlers, I.M. Belyakov. 2000. Design of Engineered Vaccines for HIV. In: Wong-Staal F, Gallo RC, editors. *AIDS Vaccine Research In Perspective*. New York: Marcel Dekker, Inc.; in press.
312. Nakagawa Y, T. Takeshita, J.A. Berzofsky, H. Takahashi. 2000. Analysis of mechanism for extracellular processing in the presentation of HIV-1 envelope protein derived peptide to epitope-specific CTL. *Immunology*. 101: 76-82..
313. Grene E, D.A. Newton, E.A. Brown, J.A. Berzofsky, S. Gattoni-Celli, G.M. Shearer. 2000. Semi-allogeneic cell hybrids stimulate HIV-1 envelope-specific cytotoxic T lymphocytes. *AIDS*: 14: 1497-1506.
314. Polakova, K., D. Plaksin, D.H. Chung, I.M. Belyakov, J.A. Berzofsky, and D.H. Margulies, 2000. Antibodies directed against the MHC-I molecule H-2D^d complexed with an antigenic peptide: similarities to a T cell receptor with the same specificity. *J. Immunol.* 165: 5703-5712.
315. Belyakov, I.M., J.D. Ahlers, J.D. Clements, W. Strober, and J.A. Berzofsky, 2000. Interplay of cytokines and adjuvants in the regulation of mucosal and systemic HIV-specific cytotoxic T lymphocytes. *J. Immunol.* 165: 6454-6462.

316. Terabe, M., S. Matsui, N. Noben-Trauth, H. Chen, C. Watson, D. D. Donaldson, D. P. Carbone, W. E. Paul, and J. A. Berzofsky. 2000. NKT cell-mediated repression of tumour immunosurveillance by IL-13 and the IL-4R-STAT6 pathway. *Nature Immunology* 1: 515-520.
317. Takahashi, M., Y. Nakagawa, J. A. Berzofsky, and H. Takahashi. 2001. Counter-regulation of cytolytic activity and cytokine production in human immunodeficiency virus (HIV)-1-specific murine CD8⁺ cytotoxic T lymphocytes by free antigenic peptide. *Int Immunol.* 13: 43-51.
318. Derby, M. A., M. A. Alexander-Miller, R. Tse, and J. A. Berzofsky. 2001. High avidity CTL exploit two complementary mechanisms to provide better protection against viral infection than low avidity CTL. *J. Immunol.*, 166: 1690-1697.
319. Shirai, M., R. Fujinaga, T. Masaki, and J. A. Berzofsky. 2001. Impaired development of HIV-1 gp160-specific CD8⁺ cytotoxic T cells by a delayed switch from Th1 to Th2 cytokine phenotype in mice with *Helicobacter pylori* infection. *Eur. J. Immunol.*, 31: 516-526.
320. Simon, R. M., S. M. Steinberg, M. Hamilton, A. Hildesheim, S. Khleif, L. W. Kwak, C. L. Mackall, J. Schlom, S. L. Topalian, and J. A. Berzofsky. 2001. Clinical trial designs for the early clinical development of therapeutic cancer vaccines. *J. Clin. Oncol.* 19: 1848-1854.
321. McGettigan, J. P., H. D. Foley, I. M. Belyakov, J. A. Berzofsky, R. J. Pomerantz, and M. J. Schnell. 2001. Rabies virus-based vectors expressing HIV-1 envelope protein induce a strong, cross-reactive cytotoxic T-lymphocyte response against envelope proteins from different HIV-1 isolates. *J. Virol.* 75: 4430-4434.
322. Derby, M.A., J. Wang, D. H. Margulies, and J. A. Berzofsky. 2001. Two intermediate avidity CTL clones with a disparity between functional avidity and MHC tetramer staining. *Internat. Immunol.* 13: 817-824.
323. Ahlers, J. D., I. M. Belyakov, S. Matsui, and J. A. Berzofsky. 2001. Mechanisms of cytokine synergy essential for vaccine protection against viral challenge. *Internat. Immunol.* 13: 897-908.
324. Wong, E. C. C., V. E. Maher, K. Hines, J. Lee, C. S. Carter, T. Goletz, W. Kopp, C. L. Mackall, J. A. Berzofsky, and E. J. Read. 2001. Development of a clinical-scale method for generation of dendritic cells from peripheral blood monocytes for use in cancer immunotherapy. *Cytotherapy* 3: 19-29.
325. Chung, D. H., I. M. Belyakov, M. A. Derby, J. Wang, L. F. Boyd, J. A. Berzofsky, and D. H. Margulies. 2001. Competitive inhibition *in vivo* and skewing of the T cell repertoire of antigen-specific CTL priming by an anti-peptide-MHC mAb. *J. Immunol.* 167: 699-707.

326. Horner, A. A., S. K. Datta, K. Takabayashi, I. M. Belyakov, T. Hayashi, N. Cinman, M.-D. Nguyen, J. H. Van Uden, J. A. Berzofsky, D D. Richman, and E. Raz. 2001. Immunostimulatory DNA-based vaccines elicit multifaceted immune responses against HIV at systemic and mucosal sites. *J. Immunol.*, in press.
327. Ahlers, J. D., I. M. Belyakov, S. Matsui, and J. A. Berzofsky. 2001. Signals delivered through TCR instruct IL-12R expression: IL-12 and TNF α synergize for IL-12R expression at low antigen dose. *Internat. Immunol.*, in press.
328. Derby, M. A., J. T. Snyder, R. Tse, M. A. Alexander-Miller, and J. A. Berzofsky. 2001. An abrupt and concordant initiation of apoptosis: antigen-dependent death of CD8⁺ CTL. *Eur. J. Immunol.*, in press.
329. Worley, B. S., L. T. van den Broeke, T. J. Goletz, C. D. Pendleton, E. M. Daschbach, E. K. Thomas, F. M. Marincola, L. J. Helman, and J. A. Berzofsky. 2001. Antigenicity of fusion proteins from sarcoma-associated chromosomal translocations. *Cancer Research* 61: in press.

330. Hel, Z., J. Nacsa, B. Kelsall, W.-P. Tsai, N. Letvin, R. W. Parks, E. Trynieszewska, L. Picker, M. G. Lewis, Y. Edghill-Smith, M. Moniuszko, R. Pal, L. Stevceva, J. D. Altman, T. M. Allen, D. Watkins, J. V. Torres, J. A. Berzofsky, I. M. Belyakov, W. Strober, and G. Franchini. 2001. Impairment of gag-specific CD8⁺ T-cell function in mucosal and systemic compartments of SIV_{mac251}- and SHIV_{ku2}- infected macaques. *J. Virol.*, in press.

Jeffrey B. Ulmer*,
John J. Donnelly* and
Margaret A. Liu

Department of Virus and Cell
Biology, Merck Research
Laboratories, West Point

Presentation of an exogenous antigen by major histocompatibility complex class I molecules

Cytotoxic T lymphocytes (CTL) generally recognize peptides derived from endogenously expressed proteins in association with nascent major histocompatibility complex (MHC) class I molecules. In contrast, peptides derived from exogenous proteins associate with MHC class II following endocytosis to an endosomal compartment. However, we have recently demonstrated that exogenous fusion proteins consisting of the binding and translocating domains of *Pseudomonas* exotoxin (PE) fused with CTL epitopes derived from either influenza matrix protein (PEMa) or nucleoprotein are internalized, processed, targeted to and presented by MHC class I (Donnelly et al. 1993, Proc. Natl. Acad. Sci. USA 1993. 90: 3530). PE is known to be internalized, processed in endosomes, and translocated to the cytosol during intoxication of cells. However, our present studies demonstrate that, unlike PE, PEMa does not require translocation to the cytosol to exert its effect. First, two inhibitors of PE toxicity that exert their effects at steps subsequent to endosomal processing had no effect on the sensitization of target cells for CTL-mediated lysis by PEMa. NH₄Cl, which inhibits PE by raising endosomal pH, and brefeldin A, which inhibits PE by disrupting the Golgi complex, did not inhibit sensitization of target cells by PEMa. Second, PEMa was capable of sensitizing for lysis T2 mutant cells, which are defective in transport of peptides from the cytosol to the lumen of the endoplasmic reticulum for presentation by MHC class I. These results suggest that PEMa is proteolytically processed in endosomes, and association with MHC class I does not require nascent MHC molecules. Such a process may involve internalized MHC class I, and subsequent expression of the peptide-MHC complexes on the cell surface would then lead to recognition by CTL.

1 Introduction

Cytotoxic T lymphocyte (CTL) responses are generated by presentation of antigens to CD8⁺ T lymphocytes by MHC class I molecules. A key to this process is the intracytosolic presence of antigens, such as in the case of endogenously synthesized viral proteins. Because CTL are important for protection against viral infections [1-4] and tumors [5] and because CTL are able to recognize epitopes derived from viral proteins, it is desirable to develop vaccines designed to engender a CTL response. Such a response has been generated *in vivo* using plasmid DNA [6], or retroviral vectors or modified pathogens [7-9] to express foreign proteins *in situ*. Alternatively, exogenous proteins and peptides have been delivered intracellularly for presentation by MHC class I, mediated by immunostimulating complexes [10], saponin adjuvants [11], synthetic lipopeptides [12], and pH-sensitive liposomes [13]. Recently, we described a recombinant protein consisting of the binding and translocating domains of *Pseudomonas* exotoxin A (PE) fused with the HLA-A2.1-restricted CTL epitope of

influenza matrix protein [14]. The native toxin is internalized by cell surface proteins, undergoes intracellular processing, and translocates its catalytic domain to the cytosol [15, 16]. The PE fusion protein (PEMa) was internalized and processed for presentation by MHC class I in target cells of the appropriate haplotype. The initial phase of intracellular trafficking of PEMa was shown to be similar to that of the toxin, in that a mutant of PEMa corresponding to a point mutation in PE that inhibits endosomal processing was inhibitory for sensitization of cells for CTL-mediated lysis by PEMa. In the present study, the intracellular trafficking of PEMa was further characterized. Surprisingly, delivery of PEMa to the cytosol was not required for presentation of the matrix epitope by MHC class I molecules. The implications of these findings are discussed in the context of current knowledge concerning antigen presentation by MHC class I.

2 Materials and methods

2.1 Materials

T2 and T1 cells were obtained from Peter Cresswell (Yale University); a mouse monoclonal antibody to HLA-A2.1 and HMY-A2 cells were obtained from Hans Zweerink (Merck Research Laboratories); U2-OS cells were purchased from ATCC; tissue culture supplies from Gibco; [³⁵S] methionine (~1200 Ci/mmol; 1 Ci = 37 GBq) and [⁵¹Cr] K(350-600 mCi/mg) from Amersham Corporation; brefeldin A (BFA) from Epicenter Technologies; PE from List Laboratories; FITC-conjugated goat anti-mouse IgG (Fc portion) and a mouse IgG2a control serum were from Cappel-Organon Technika.

[1 12654]

* These authors contributed equally to this work.

Correspondence: Margaret A. Liu, Department of Virus and Cell Biology, Merck Research Laboratories, Building 16-101 West Point, PA 19486, USA (Fax: 215-652-7320)

Abbreviations: BFA: Brefeldin A ER: Endoplasmic reticulum
PE: *Pseudomonas* exotoxin A PEMa: PE-influenza matrix epitope fusion protein

Key words: Antigen presentation / Cytotoxic T lymphocytes / Influenza

2.2 Cell culture

U2-OS cells were maintained in McCoy's 5A medium with 15% fetal bovine serum (FBS) and were removed from flasks with 0.1 M EDTA in phosphate-buffered saline (PBS). Human CTL specific for matrix protein were obtained from a normal HLA-A2.1⁺ donor and were maintained *in vitro* as described by Bednarek et al. [17] with minor modifications [14]. T2, T1 and HMY-A2 cells were grown in suspension in RPMI 1640 containing 10% FBS.

2.3 Biosynthetic labeling

For measurement of PE toxicity, U2-OS cells were plated in a 96-well plate at a density of 20 000 cells/well and allowed to attach overnight. Cell monolayers were treated with or without BFA or NH_4Cl for 1 h at 37°C, followed by the addition of PE (0-1000 pM final concentration) for 15 min. Cells were washed with culture medium containing BFA or NH_4Cl and incubated with [^{35}S]methionine (2 μCi) and BFA or NH_4Cl for 5 h. Cells were washed with ice-cold culture medium, lysed with sodium dodecylsulfate (0.1%) and added to an equal volume of ice-cold trichloroacetic acid (20%). Incorporation of radioactivity into acid precipitable material was determined by liquid scintillation spectrometry. For measurement of protein secretion, U2-OS cell monolayers were treated with BFA (1-10 $\mu\text{g}/\text{ml}$) for 30 min at 37°C, followed by the addition of [^{35}S]methionine. Aliquots of cell culture supernatant were taken at various times and spotted onto filter papers. Secreted proteins were precipitated onto the filters by soaking in ice-cold trichloroacetic acid (10%) for 1 h. Filters were washed three times in ice-cold trichloroacetic acid (5%) and three times in ice-cold ethanol, dried and radioactivity was determined as before.

2.4 Cytotoxic T lymphocyte assay

U2-OS cells were plated in 24-well plates at 4×10^5 cells per well and labeled overnight with 2 $\mu\text{Ci}/\text{ml}$ of ^{51}Cr . The following day, cell monolayers or suspensions were exposed to BFA (1-10 $\mu\text{g}/\text{ml}$) or NH_4Cl (10-80 mM) for designated periods of time. Cells were incubated with PEMa, washed, and in some cases further incubated in the absence of PEMa, and finally washed with PBS. For BFA-treated target cells, all wash solutions contained BFA. To obtain influenza virus-infected targets, cells were incubated with influenza A virus (A/Victoria/73) for 30 min, washed, and incubated for a further 2 h. EDTA (0.1 M in PBS) was then used to detach U2-OS cells from the plates, and the cells were washed twice more and resuspended in RPMI 1640 medium with 10% FBS and 0.01 M HEPES at 5×10^4 cells/ml. T1 and T2 cells were labeled overnight with 2 $\mu\text{Ci}/\text{ml}$ of ^{51}Cr , and washed by sedimentation and resuspension. Where indicated, cells were fixed with glutaraldehyde (0.1%) for 1 min on ice. Target cells were plated in triplicate at 0.1 ml per well in round-bottom, 96-well plates with 0.1 ml CTL at effector:target ratios of 30:1, 20:1 or 10:1. Cells were incubated at 37°C, sedimented, and 20 μl of supernatant was removed for counting in an LKB 1212 plate scintillation counter. Results were expressed as specific lysis = [(counts released with CTL)-(counts

released without CTL)/(counts released by 6 M HCl)-(counts released without CTL)] $\times 100$.

In bystander killing experiments, T1, T2 and HMY-A2 cells were incubated with PEMa (1 $\mu\text{g}/\text{ml}$) for 30 min, washed and added to adherent U2-OS cells that were loaded overnight with ^{51}Cr , as before. The cell mixtures were incubated for 2 h and the PEMa-pulsed T1, T2 or HMY-A2 cells were decanted. The U2-OS cells were washed and incubated with influenza-specific CTL for 2 h. In the converse experiment, U2-OS cells were incubated with PEMa, and T1, T2 or HMY-A2 cells were loaded with ^{51}Cr , then the cell mixtures were incubated as before. The T1, T2 and HMY-A2 cells were decanted and incubated with CTL for 2 h. In each case, negative controls consisted of cells without prior incubation with PEMa, as well as no addition of CTL to bystander cells. As positive controls, each set of bystander cells was incubated with PEMa or matrix peptide (57-68) to demonstrate sensitization.

2.5 Immunofluorescence staining

U2-OS cells were treated with or without BFA (1-10 $\mu\text{g}/\text{ml}$) for various periods of time (0-10 h), and removed from culture flasks with EDTA. For staining of surface MHC class I molecules, cells were washed with PBS containing FBS (5%) and sodium azide (0.01%), then incubated with a monoclonal HLA-A2.1 antibody or a normal mouse IgG2a control (12.5 $\mu\text{g}/\text{ml}$ final concentration) for 30 min on ice. Cells were washed, incubated with a FITC-conjugated goat anti-mouse IgG (Fc portion) (25 $\mu\text{g}/\text{ml}$) for 30 min on ice, and washed again. Measurements were as mean channel fluorescence intensity using a Becton Dickinson FACScan flow cytometer with an excitation wavelength of 488 nm and a band pass filter of 530 nm. A minimum of 3500 events/sample were analyzed.

3 Results

The fusion protein PEMa has been previously shown to be internalized by U2-OS and HMY A2 target cells and presented by MHC class I [14]. These cells were chosen because they express the HLA-A2.1 MHC haplotype, in order that the HLA-A2.1-restricted CTL epitope of influenza matrix protein (amino acids 57-68) would be presented by MHC class I and specifically recognized by appropriate CTL.

Additionally, these cells bind and process PE, and are sensitive to the toxin (unpublished observations). It was demonstrated that, as in the case of PE, delivery to endosomes and proteolytic processing of PEMa therein was a necessary step in its action. In the present study, we sought to characterize further the intracellular trafficking of PEMa in order to investigate the location of association with MHC class I during sensitization of target cells for lysis by CTL. This was accomplished using compounds that have been shown to block different steps in the processing of PE.

PE intoxication is known to be inhibited by weak amines, such as ammonium chloride, at a step subsequent to endosomal proteolytic processing [16]. The precise mechanism of inhibition by primary amines has not yet been

defined. But, given that PE may translocate from some other compartment such as the ER [18, 19], there are several potential events following endosomal processing that could be affected (e.g., budding of vesicles from endosomes or Golgi, binding to the KDEL receptor, or translocation to the cytosol). U2-OS cells were completely insensitive to the toxin in the presence of ammonium chloride at concentrations shown to inhibit PE in other cell types (Fig. 1 A). Under these conditions, however, sensitization of cells by PEMa for CTL-mediated lysis was not affected (Fig. 1 B). These results indicate that an obligatory step subsequent to endosomal processing of the toxin is not necessary for sensitization by PEMa and represents a divergence in the pathways of the two molecules.

Recently, BFA was shown to inhibit intoxication of cells by PE [20, 21]. Although the means of this inhibition is not fully understood, BFA treatment is known to cause: (i) a cessation in protein exocytosis due to the disruption and disappearance of the Golgi complex [22, 23], and (ii) morphologic but not functional changes in endosomes [24]. Due to the presence of a KDEL-like sequence on the C-terminus of PE that is necessary for PE toxicity [18], it

has been proposed that prior to translocation to the cytosol PE is transported from endosomes to the Golgi complex and endoplasmic reticulum (ER) in a process that involves the KDEL receptor [19]. Therefore, inhibition of PE toxicity by BFA is likely to be due to its effects on the Golgi complex and, hence, may prevent trafficking of PE to the ER for translocation to the cytosol. Treatment of U2-OS cells with BFA resulted in an inhibition of protein secretion (Fig. 2 A), and also prevented intoxication of the cells by PE (Fig. 2 B). Incubation of the cells with BFA did not have a discernible effect on sensitization by PEMa (Fig. 2 C), demonstrating that PEMa does not require delivery to, or passage through the Golgi complex, for association with MHC class I. As a positive control for the inhibitory effect of BFA on antigen presentation by nascent MHC class I, virus-infected U2-OS cells were tested. Virus-infected cells were sensitized for lysis by CTL, but not when the cells were treated with BFA following a 30-min pulse with virus (Fig. 2 D). Since evidence indicates that association of




Figure 1. Effect of ammonium chloride on PE toxicity and sensitization by PEMa. Panel A: PE toxicity. U2-OS cells were incubated in the presence (open circles) or absence (closed circles) of NH_4Cl (30 mM) and PE toxicity was measured as the inhibition of incorporation of $[^{35}\text{S}]$ methionine, as detailed in Sect. 2.3. The data are plotted as incorporation of radioactivity into acid-precipitable material, as a percentage of maximal incorporation, versus PE concentration. Error bars represent \pm SD, where $n = 6$. Panel B: Sensitization by PEMa. Cells were incubated with NH_4Cl at the indicated concentrations for 2 h at 37°C , followed by the addition of PEMa (open circles) at $1 \mu\text{M}$ for 2 h. Controls were incubated in the absence of PEMa (closed circles). Cells were fixed and incubated with CTL (effector : target = 20 : 1) for 2 h at 37°C . CTL-mediated lysis of target cells is shown as % specific lysis versus NH_4Cl concentration. Error bars represent \pm SD, where $n = 3$.




Figure 2. Effect of BFA on U2-OS cells. Panel A: Protein secretion. Cells were incubated with (open circles) or without (closed circles) BFA ($1 \mu\text{g}/\text{ml}$) and protein secretion into the culture supernatant was measured. Shown is acid-precipitable radioactivity versus time in BFA. Similar results were obtained with $10 \mu\text{g}$ BFA (not shown). Error bars represent \pm SD, where $n = 3$. Panel B: PE toxicity. Cells were incubated with (open circles) or without (closed circles) BFA ($1 \mu\text{g}/\text{ml}$) and effect of PE on protein synthesis was measured as incorporation of $[^{35}\text{S}]$ methionine. Shown is acid-precipitable radioactivity in the cells, as a percentage of maximal incorporation, versus PE concentration. Error bars represent \pm SD, where $n = 6$. Panel C: Sensitization by PEMa. Cells were incubated with (open circles) or without (closed circles) BFA ($2 \mu\text{g}/\text{ml}$) for 2 h at 37°C , followed by the addition of PEMa ($1 \mu\text{M}$) for 4 h. The cells were washed with culture medium containing BFA and incubated with CTL at an effector : target ratio of 20 : 1 for an additional 1.5 h in the presence of BFA. At this concentration, BFA had no detrimental effect on the ability of CTL to lyse target cells (not shown). Results are plotted as % specific lysis versus PEMa concentration. Error bars represent \pm SD, where $n = 3$. Panel D: Sensitization by virus infection. Cells were: (i) untreated, (ii) incubated with PEMa ($10 \mu\text{M}$) in the presence or absence of BFA, or (iii) incubated with influenza A virus (A/Victoria/73), for 30 min and washed. The cells were incubated a further 2 h in the presence (striped bars) or absence (solid bars) of BFA ($2 \mu\text{g}/\text{ml}$) and finally with CTL (25 : 1 E : T) for 2 h. Error bars represent \pm SD, where $n = 3$.

peptides with nascent MHC class I occurs in the ER [25, 26], it is unlikely that the influenza matrix epitope from PEMA associated with nascent MHC class I. It is possible, though, that internalized PEMA could have intersected with nascent MHC class I in a compartment that is spared by BFA treatment, such as the *trans* Golgi network (TGN), during transit to the cell surface. However, it was recently shown that BFA inhibits the constitutive and regulated secretion of proteins from the TGN to the plasmalemma [27].

Therefore, MHC class I and influenza matrix peptides, if present together in the TGN during BFA treatment, would be trapped and not reach the cell surface. Moreover, sensitization of cells by PEMA was observed after prolonged treatment of cells with BFA (10 h) (data not shown), which based on intracellular transport of MHC class I molecules in other cells [28, 29] should have been sufficient time to clear the transport vesicles containing newly synthesized MHC class I operating between the TGN and cell surface. Therefore, it is more likely that association of the matrix epitope derived from PEMA occurred in an intracellular compartment containing preexisting MHC class I molecules. It is perhaps not surprising that BFA did not inhibit sensitization by PEMA since PE fusion proteins were capable of sensitizing target cells for lysis whether or not the KDEL-like sequence was present on the C-terminus [14]. Sensitization by a PE fusion protein lacking a KDEL-like sequence suggests that transport to the ER by the KDEL receptor is not required for delivery to MHC class I and further supports the conclusion that nascent MHC class I molecules are not involved.

To test this hypothesis further, T2 mutant cells were used. These cells contain a deletion in the MHC region of the genome [30, 31] and are defective in the transport of peptides from the cytosol to the lumen of the ER for association with MHC class I [32]. Yet, these cells express HLA-A2 molecules on the cell surface due to presentation of peptides derived from the signal sequences of endogenous proteins [33]. Therefore, if delivery of PEMA to the cytosol is a necessary step prior to association with MHC class I, PEMA would not be capable of sensitizing T2 cells for lysis by CTL. In fact, PEMA sensitized T2 cells with the same or better efficiency as wild-type T1 cells (Fig. 3).




Figure 3. Sensitization of T2 cells by PEMA. T1 (closed circles) and T2 (open circles) cells were incubated with PEMA for 15 min at 37°C, washed and incubated for a further 3 h. CTL were added at an effector:target ratio of 10:1 for 3 h at 37°C. Shown is % specific lysis of target cells versus PEMA concentration. Error bars represent \pm SD, where $n = 3$.




Figure 4. Lack of sensitization of U2-OS cells for lysis by excretion of peptide by antigen presenting cells. U2-OS cells served as bystander cells and were loaded with ^{51}Cr overnight. T1, T2 and HMY-A2 cells served as antigen presenting cells and were incubated with (+) or without (-) PEMA (1 $\mu\text{g}/\text{ml}$) for 30 min and washed. The different cell mixtures were incubated together, separated and further incubated with influenza-specific CTL as described in Sect. 2.4. As positive controls, U2-OS cells were incubated with PEMA or matrix peptide (1 $\mu\text{g}/\text{ml}$) for 2 h, then incubated with CTL. Error bars represent \pm SD, where $n = 3$.

These results suggest that PEMA is internalized to an intracellular compartment other than the cytosol, Golgi complex or ER where it associates with MHC class I for eventual presentation on the cell surface.

To address the possibility that sensitization of target cells with PEMA was accomplished by internalization and processing of PEMA followed by excretion of peptide for association with cell surface MHC class I, bystander killing experiments were performed. In one set of experiments, U2-OS cells were used as bystanders and MHC haplotype-matched T1, T2 or HMY-A2 cells served as antigen presenting cells pulsed with PEMA (Fig. 4). In the converse experiment, T1, T2 and HMY-A2 cells were used as bystanders and U2-OS cells were PEMA-pulsed (Fig. 5). Sensitization of bystander cells by excretion of peptide from PEMA-pulsed cells was not detected with any combination of cells. As positive controls, each bystander cell type was incubated with PEMA or peptide in the absence of other cells and shown to be sensitized for lysis by CTL. Therefore, it is unlikely that excretion of peptides accounts for the observed sensitization by PEMA.

To investigate potential internalization of MHC class I in U2-OS cells, surface levels were analyzed by immunofluorescence staining in the presence or absence of BFA. During the period of BFA treatment, cell number remained constant and cell viability, as measured by trypan blue exclusion, was not different from that of control cells (data not shown). After 10 h in BFA, cell surface MHC class I levels decreased to approximately 85% of the starting amount (Fig. 6A, B). Samples collected at intermediate time points showed that MHC class I molecules disappeared from the cell surface at a constant rate of $\sim 1.5\%/h$ (Fig. 6C). This suggests that MHC class I underwent internalization, denaturation, and/or shedding from the cell surface with a half-time of approximately 33 h, which is comparable to that in other cells [34].




Figure 5. Lack of sensitization of T1, T2 and HMY-A2 cells by excretion of peptide from PEMA-pulsed U2-OS cells. T1 (panel A), T2 (panel B) and HMY-A2 (panel C) cells served as bystanders and were incubated with ^{51}Cr overnight. U2-OS cells served as antigen presenting cells and were incubated with (+) or without (-) PEMA (1 $\mu\text{g}/\text{ml}$). Bystanders and PEMA-pulsed cells were incubated together and treated as described in Fig. 4. As positive controls, each bystander cell type was incubated with PEMA or peptide. Error bars represent \pm SD, where $n = 3$.

Taken together, these results: (i) demonstrate that intracellular trafficking of PEMA contrasts that of PE, in that sensitization of cells by PEMA does not require delivery from endosomes to the Golgi complex, ER or cytosol, (ii) demonstrate that sensitization by PEMA does not involve excretion of peptide for association with cell surface MHC class I, and (iii) suggest that U2-OS cell surface MHC class I molecules may undergo constitutive internalization.

4 Discussion

The nature of antigen presentation by MHC class I has been well studied and it is clear that nascent MHC molecules in the ER associate with peptides derived from

endogenously synthesized antigens, such as viral proteins [25, 26]. However, it has been observed that, by using various vehicles, exogenous antigens can be delivered to the cytosol and presented by MHC class I molecules [10-13, 35, 36]. Exogenous proteins by themselves are generally not able to enter an MHC class I processing pathway. But, by utilizing fusion proteins consisting of the binding and translocating domains of PE, which has intrinsic capabilities to gain entry into intracellular compartments, we have succeeded in delivering exogenous proteins containing CTL epitopes to such a pathway. The key findings of this study are as follows: (i) the intracellular pathway taken by PEMA differs from that of PE at a point at or past endosomes, (ii) sensitization of target cells by PEMA does not require delivery of the matrix epitope to the cytosol, and (iii) association of the matrix epitope with MHC class I does not require newly synthesized MHC molecules. One might predict that delivery of PEMA to nascent MHC molecules in the ER would be inefficient, due to the lack of a KDEL-like sequence on the C-terminus of the fusion protein. Such a sequence on PE was recently




Figure 6. Cell surface staining of MHC class I. Cells were incubated with BFA (1 $\mu\text{g}/\text{ml}$) for up to 10 h and aliquots of cells were taken for immunofluorescence determination of cell surface MHC class I levels. Relative fluorescence intensity is plotted on the abscissa versus frequency on the ordinate for cells treated with BFA for 10 h (open peak) and untreated control (closed peak). Staining with monoclonal HLA-A2.1 antibody is shown in panel A, and with isotype-matched control antibody in panel B. MHC class I levels did not change during this time period in untreated cells (not shown). Samples taken at the intermediate time points of 2, 4, 6 and 8 h were also analyzed and are plotted together with the 0 and 10 h time points as percent maximal fluorescence intensity versus time (panel C).

shown to be obligatory for intoxication [18] and implies the involvement of the KDEL receptor, which is localized in the ER and Golgi complex [37], in the delivery of the toxin to the ER for translocation to the cytosol [19].

Taken together, the results in this report suggest that the matrix epitope was able to associate with preexisting MHC class I molecules in an endocytic compartment, such as could occur by the internalization of cell surface MHC class I. The generation of peptides for association with MHC class I could be the result of proteolytic processing of PEMA in endosomes. In support of this possibility, we have isolated fragments of PEMA from endosomes of cells incubated with radiolabelled PEMA (unpublished observations). Subsequent presentation of the matrix epitope in association with MHC class I could conceivably occur by recycling of the complex to the cell surface. These possible events remain to be established; however several published reports support the hypothesis. Endocytosis and recycling of MHC class I molecules to the cell surface has been described in activated T lymphocytes [38] and T cell lines [39, 40], and may be involved in antigen presentation [41], but it is not certain whether this can be generalized to other cell types as an ongoing process.

Internalization of MHC class I molecules has been reported to occur constitutively in T lymphocytes [42], macrophage/monocyte cells [43] and T and B lymphoid cell lines [44], and can be induced in fibroblasts by cross-linking [45]. Regulated internalization was also suggested in some cases, as evidenced by phorbol ester-induced enhancement of uptake [43, 44]. Incubation of P815 cells with a photoaffinity-labeled CTL epitope resulted in specific binding to MHC class I, some of which was subsequently localized intracellularly [46], suggesting that MHC class I molecules were spontaneously internalized. In addition, it has been proposed that the virus-induced disappearance of cell surface MHC class I molecules in BGIMK primate cells is caused by an impairment of normal MHC class I recycling [47]. Finally, two recent reports suggest the involvement of endolysosomal processing of antigens for presentation by MHC class I. First, CTL-mediated lysis of Sendai virus-infected T2 cells was not inhibited by BFA treatment [48], indicating that an alternative pathway can function in the presentation of antigens by MHC class I. Second, bacterial antigens were internalized, processed in a phagocytic compartment and presented by MHC class I molecules in BFA- and cycloheximide-treated cells [49]. In that study, it was postulated that intracellular antigen processing and excretion of peptide led to the expression of peptide-MHC complexes on the cell surface. However, the results could also be explained by internalization and recycling of cell surface MHC class I. In the present study, excretion of the matrix epitope by PEMA-pulsed cells was not detected, as measured by the lack of sensitization of haplotype-matched bystander cells.

In contrast to the cited examples of MHC class I internalization and recycling, internalization of MHC class I molecules in the JY human B lymphoblastoid cell line [50, 51] and in B lymphocytes [42] was not detected. Therefore, this process may depend on cell type or may be more prominent in some cells than others (for review see [52]). The results presented in this report, though, suggest that MHC class I molecules may not only be internalized to an endocytic

compartment but also may associate with antigenic peptides therein and recycle back to the plasma membrane in non-lymphoid cells, where they can be recognized by appropriate CTL. Such a pathway could explain why certain exogenously applied proteins without an apparent means of gaining access to the cytosol are capable of being presented by MHC class I molecules *in vitro* and/or inducing a CTL response *in vivo* [53-59]. That exogenous proteins in general do not enter this processing pathway suggests there is limited access. However, the use of ligands, such as PE, to target exogenous antigens containing CTL epitopes for endocytosis to intracellular processing compartments may facilitate their presentation by MHC class I, and may form the basis of vaccine efforts designed to generate CTL responses. To provide determinant selection in an outbred population, however, it will be necessary to first determine whether larger polypeptides can be targeted to cells and processed for presentation by MHC class I using this technique. Nevertheless, because of the sensitivity of CTL in recognizing antigen/MHC class I complexes [60], presentation of exogenous toxin/antigen fusion proteins by a putative recycling MHC class I pathway has potential as a means of eliciting CTL responses *in vivo*.

We thank Xiao-Ping Shi, Donna Montgomery and Karen Leander (Merck Research Laboratories) for molecular biology support; Arthur Friedman and Douglas Martinez (Merck Research Laboratories) for protein purification; Linda Hawe for help with CTL assays; Chenyang Shi (National University of Singapore) for assistance with the immunofluorescence staining studies; and Peter Cresswell (Yale) and Hans Zveierink (Merck Research Laboratories) for reagents.

Received January 6, 1994; in final revised form April 19, 1994; accepted April 20, 1994.

5 References

- 1 Rouse, B. T., Norley, S. and Martin, S., *Rev. Infect. Dis.* 1988. 10: 16.
- 2 Lukacher, A. E., Braciale, V. L. and Braciale, T. J., *J. Exp. Med.* 1984. 160: 814.
- 3 McMichael, A. J., Gotch, F. M., Noble, G. R. and Beare, P. A. S., *N. Engl. J. Med.* 1983. 309: 13.
- 4 Wraith, D. C., Vessey, A. E. and Askonas, B. A., *J. Gen. Virol.* 1987. 68: 433.
- 5 Cerundolo, V., Lahaye, T., Horvath, C., Zanovello, P., Collavo, D. and Engers, H. E., *Eur. J. Immunol.* 1987. 17: 173.
- 6 Ulmer, J. B., Donnelly, J. J., Parker, S. E., Rhodes, G. H., Felgner, P. L., Dwarki, V. J., Gromkowski, S. H., Deck, R. R., DeWitt, C. M., Friedman, A., Hawe, L. A., Leander, K. R., Martinez, D., Perry, H. C., Shiver, J. W., Montgomery, D. L. and Liu, M. A., *Science* 1993. 259: 1745.
- 7 Schafer, R., Portnoy, D. A., Brassel, S. A. and Paterson, Y., *J. Immunol.* 1992. 149: 53.
- 8 Hahn, C. S., Hahn, Y. S., Braciale, T. J. and Rice, C. M., *Proc. Natl. Acad. Sci. USA* 1992. 89: 2679.
- 9 Aldovini, A. and Young, R. A., *Nature* 1992. 351: 479.
- 10 Takahashi, H., Takeshita, T., Morein, B., Putney, S., Germain, R. N. and Berzofsky, J. A., *Nature* 1990. 344: 873.
- 11 Newman, M. J., Wu, J.-Y., Gardner, B. H., Munroe, K. J., Leombruno, D., Recchia, J., Kensil, C. R. and Coughlin, R. T., *J. Immunol.* 1992. 148: 2357.
- 12 Deres, K., Schild, H., Weismüller, K.-H., Jung, G. and Rammensee, H. G., *Nature* 1992. 342: 561.

- 13 Collins, D. S., Findlay, K. and Harding, C. V., *J. Immunol.* 1992. 148: 3336.
- 14 Donnelly, J. J., Ulmer, J. B., Hawe, L. A., Friedman, A., Shi, X.-P., Leander, K. R., Shiver, J. W., Oliff, A. I., Martinez, D., Montgomery, D. and Liu, M. A., *Proc. Natl. Acad. Sci. USA* 1993. 90: 3530.
- 15 Kuonnas, M. Z., Morris, R. E., Thompson, M. R., FitzGerald, D. J., Strickland, D. K. and Saelinger, C. B., *J. Biol. Chem.* 1992. 267: 12420.
- 16 Ogata, M., Chaudhary, V. K., Pastan, I. and FitzGerald, D., *J. Biol. Chem.* 1990. 265: 20678.
- 17 Bednarek, M. A., Engl, S. A., Gammon, M. C., Linquist, J. A., Porter, G., Williamson, A. R. and Zweerink, H. J., *J. Immunol. Methods* 1991. 139: 41.
- 18 Chaudhary, V. K., Jinno, Y., FitzGerald, D. and Pastan, I., *Proc. Natl. Acad. Sci. USA* 1990. 87: 308.
- 19 Pelham, H. R. B., Roberts, L. M. and Lord, J. M., *Trends Cell Biol.* 1992. 2: 183.
- 20 Yoshida, T., Chen, C., Zhang, M. and Wu, H. C., *Exp. Cell Res.* 1991. 192: 389.
- 21 Seetharam, S., Chaudhary, V. K., FitzGerald, D. and Pastan, I., *J. Biol. Chem.* 1991. 266: 17376.
- 22 Misumi, Y., Misumi, Y., Miki, K., Takatsuki, A., Tamura, G. and Ikehara, Y., *J. Biol. Chem.* 1986. 261: 11398.
- 23 Ulmer, J. B. and Palade, G. E., *Eur. J. Cell Biol.* 1991. 54: 38.
- 24 Wood, S. A. and Brown, W. J., *J. Cell Biol.* 1992. 119: 273.
- 25 Nuchtern, J. G., Bonifacio, J. S., Biddison, W. E. and Klausner, R. D., *Nature* 1989. 339: 223.
- 26 Cox, J. H., Yewdell, J. W., Eisenlohr, L. C., Johnson, P. R. and Bennink, J. R., *Science* 1990. 247: 715.
- 27 Rosa, P., Barr, F. A., Stinchcombe, J. C., Binacchi, C. and Huttner, W. B., *Eur. J. Cell Biol.* 1992. 59: 265.
- 28 Williams, D. B., Swiedler, S. J. and Hart, G. W., *J. Cell Biol.* 1985. 101: 725.
- 29 Williams, D. B., Borriello, F., Zeff, R. A. and Nathenson, S. G., *J. Biol. Chem.* 1988. 263: 4549.
- 30 DeMars, R., Rudersdorf, R., Chang, C., Petersen, J., Strandtmann, J., Korn, N., Sidwell, B. and Orr, H. T., *Proc. Natl. Acad. Sci. USA* 1985. 82: 8183.
- 31 Salter, R. D., Howell, D. N. and Cresswell, P., *Immunogenetics* 1985. 21: 235.
- 32 Anderson, K., Cresswell, P., Gammon, M., Hermes, J., Williamson, A. and Zweerink, H., *J. Exp. Med.* 1991. 174: 489.
- 33 Wei, M. L. and Cresswell, P., *Nature* 1992. 356: 443.
- 34 Male, D. and Pryce, G., *Immunology* 1988. 63: 37.
- 35 Harding, C. V., *Eur. J. Immunol.* 1992. 22: 1865.
- 36 Yewdell, J. W., Bennink, J. R. and Hosaka, Y., *Science* 1988. 239: 637.
- 37 Lewis, M. J. and Pelham, H. R. B., *Nature* 1990. 348: 162.
- 38 Tse, D. B. and Pernis, B., *J. Exp. Med.* 1984. 159: 193.
- 39 Reid, P. A. and Watts, C., *Nature* 1990. 346: 655.
- 40 Vega, M. A. and Strominger, J. L., *Proc. Natl. Acad. Sci. USA* 1989. 86: 2688.
- 41 Hochman, J. H., Jiang, H., Matyus, L., Edidin, M. and Pernis, B., *J. Immunol.* 1991. 146: 1862.
- 42 Machy, P., Truneh, A., Gennaro, D. and Hoffstein, S., *Nature* 1987. 328: 724.
- 43 Dasgupta, J. D., Watkins, S., Slayter, H. and Yunis, E. J., *J. Immunol.* 1988. 141: 2577.
- 44 Capps, C. G., Van Kampen, M., Ward, C. L. and Zúñiga, M. C., *J. Cell Biol.* 1989. 108: 1317.
- 45 Huet, C., Ash, J. F. and Singer, S. J., *Cell* 1980. 21: 429.
- 46 Cerottini, J.-C. and Luescher, I. F., *Immunol. Lett.* 1991. 30: 171.
- 47 Boshkov, L. K., Macen, J. L. and McFadden, G., *J. Immunol.* 1992. 148: 881.
- 48 Zhou, X., Glas, R., Ljunggren, H.-G. and Jondal, M., *Eur. J. Immunol.* 1993. 23: 1802.
- 49 Pfeifer, J. D., Wick, M. J., Roberts, R. L., Findlay, K., Normark, S. J. and Harding, C. V., *Nature* 1993. 361: 359.
- 50 Neeffjes, J. J., Stollorz, V., Peters, P. J., Geuze, H. J. and Ploegh, H. L., *Cell* 1990. 61: 171.
- 51 Eichholtz, T., Vossebeld, P., van Overveld, M. and Ploegh, H., *J. Biol. Chem.* 1992. 267: 22490.
- 52 Pernis, B., *Immunol. Today* 1985. 6: 45.
- 53 Tevethia, S. S., Flyer, D. C. and Tjian, R., *Virology* 1980. 107: 13.
- 54 Yamada, A., Ziese, M. R., Young, J. E., Yamada, Y. K. and Ennis, F. A., *J. Exp. Med.* 1985. 162: 663.
- 55 Wraith, D. C. and Vessey, A. E., *Immunology* 1986. 59: 173.
- 56 Staerz, U. E., Karasuyama, H. and Garner, A. M., *Nature* 1987. 329: 449.
- 57 Jin, Y., Shih, J. W.-K. and Berkower, I., *J. Exp. Med.* 1988. 168: 293.
- 58 Carbone, F. R. and Bevan, M. J., *J. Exp. Med.* 1990. 171: 377.
- 59 Rock, K. L., Gamble, S. and Rothstein, L., *Science* 1990. 249: 918.
- 60 Christnick, E. R., Luscher, M. A., Barber, B. H. and Williams, D. B., *Nature* 1991. 352: 67.

Targeting HIV proteins to the major histocompatibility complex class I processing pathway with a novel gp120-anthrax toxin fusion protein

THERESA J. GOLETZ^{*†}, KURT R. KLIMPEL^{†‡}, NAVEEN ARORA[‡], STEPHEN H. LEPPLA^{‡§}, JERRY M. KEITH[‡],
AND JAY A. BERZOFKY^{*}

^{*}Molecular Immunogenetics and Vaccine Research Section, Metabolism Branch, Division of Clinical Sciences, National Cancer Institute, Bethesda, MD 20892; and the [‡]Oral Infection and Immunity Branch, National Institute of Dental Research, Bethesda, MD 20892

Communicated by John B. Robbins, National Institute of Child Health and Development, Bethesda, MD, August 25, 1997 (received for review April 8, 1997)

ABSTRACT A challenge for subunit vaccines whose goal is to elicit CD8⁺ cytotoxic T lymphocytes (CTLs) is to deliver the antigen to the cytosol of the living cell, where it can be processed for presentation by major histocompatibility complex (MHC) class I molecules. Several bacterial toxins have evolved to efficiently deliver catalytic protein moieties to the cytosol of eukaryotic cells. Anthrax lethal toxin consists of two distinct proteins that combine to form the active toxin. Protective antigen (PA) binds to cells and is instrumental in delivering lethal factor (LF) to the cell cytosol. To test whether the lethal factor protein could be exploited for delivery of exogenous proteins to the MHC class I processing pathway, we constructed a genetic fusion between the amino-terminal 254 aa of LF and the gp120 portion of the HIV-1 envelope protein. Cells treated with this fusion protein (LF254-gp120) in the presence of PA effectively processed gp120 and presented an epitope recognized by HIV-1 gp120 V3-specific CTL. In contrast, when cells were treated with the LF254-gp120 fusion protein and a mutant PA protein defective for translocation, the cells were not able to present the epitope and were not lysed by the specific CTL. The entry into the cytosol and dependence on the classical cytosolic MHC class I pathway were confirmed by showing that antigen presentation by PA + LF254-gp120 was blocked by the proteasome inhibitor lactacystin. These data demonstrate the ability of the LF amino-terminal fragment to deliver antigens to the MHC class I pathway and provide the basis for the development of novel T cell vaccines.

Whereas serum IgG are believed to be sufficient for protection conferred by most current vaccines (1), CD8⁺ cytotoxic T cells (CTLs) are believed to be important in clearing viral infections (1–8). A more vigorous CTL response against a viral infection thus should be beneficial (9–13). Thus, vaccines eliciting specific CTLs should be valuable, especially in the case of viruses that produce chronic infections, such as HIV. Although live viruses elicit CTL immunity, soluble whole proteins, such as subunit vaccines, with or without adjuvants, generally do not (14–17). Because whole exogenous proteins are usually internalized via the endosomal system, they are processed for presentation by class II major histocompatibility complex (MHC) molecules, generally not class I. In contrast, presentation by class I MHC molecules typically occurs only for proteins proteolytically processed in the cytoplasm of the cell. Cytosolic proteins are digested into short peptides by proteasomes, and the resulting peptides are transported by transporters of antigenic peptides into the endoplasmic reticulum,

where they bind to nascent class I MHC molecules, which carry them out to the cell surface for display to CD8⁺ T cells (17). Therefore, CD8⁺ CTLs, which uniquely recognize peptide epitopes presented by class I MHC molecules, provide surveillance against abnormal proteins synthesized in the cell, such as viral proteins or tumor antigens. The challenge in designing a nonliving vaccine for induction of CD8⁺ CTLs thus is to introduce the vaccine protein into the cytosol of cells for processing and presentation by their class I molecules.

Several CTL vaccine strategies have been employed with varying effectiveness, including live attenuated bacterial and viral strains, immunostimulatory complexes (ISCOMs) and liposome delivery systems for soluble antigens, and DNA-based vaccines. Unfortunately, each of these systems contains its own unique set of problems and is not always suitable for all individuals (18–21). Particularly, in the case of HIV-1 and its macaque close cousin simian immunodeficiency virus (SIV), live attenuated virus has been shown to protect adult monkeys from a subsequent challenge with virulent virus (22), but the safety of an attenuated strain, especially in newborns, has been questioned (23).

A number of bacterial toxins contain catalytic domains that share the ability to gain access to the host cell cytosol, where they can exert their effects. Although each toxin may differ in the mechanism or route by which it gains entry to the cytosol, the overall effect is that of a “molecular syringe” that is able to inject the toxic protein into the cell. Several bacterial toxins, including diphtheria toxin (DT), *Pseudomonas* exotoxin A (PE), pertussis toxin, and the pertussis adenylate cyclase, have been used in attempts to deliver peptide epitopes to the cell cytosol as internal or amino-terminal fusions (24–27). These systems are restricted in their use as potential vaccines, because their capacity to deliver larger protein antigens is limited and many individuals have already been immunized against the carrier toxin.

Although peptides are able to stimulate a cellular immune response, whole protein antigens may be better suited for use in an effective vaccine for two reasons. First, the epitope that is essential for protection in one genetic background may prove to be irrelevant in a different genetic background. Therefore, it is beneficial for a broadly applied T cell vaccine to use the full-length protein from which the various relevant epitopes are derived. Second, peptides recognized by CTLs are processed from the whole protein by specialized degradative machinery, including the proteasome complex. In certain

Abbreviations: LF, lethal factor; PA, protective antigen; CTL, cytotoxic T lymphocyte; MHC, major histocompatibility complex; PE, *Pseudomonas* exotoxin A; GST, glutathione S-transferase.

[†]T.J.G. and K.R.K. contributed equally to this work.

[§]To whom reprint requests should be addressed at: Oral Infection and Immunity Branch, National Institute of Dental Research, Bldg. 30, Rm. 309, National Institutes of Health, Bethesda, MD 20892.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.

0027-8424/97/9412059-06\$05.00/0

PNAS is available online at <http://www.pnas.org>.

instances, the processing of the relevant peptide epitopes is dependent on the flanking amino acid sequences (28). However, flanking residues are not always important for proper processing (29). Because it currently is not possible to accurately predict which epitopes are dependent on their context for proper processing, it is important to deliver the entire antigen to the cell cytosol for optimal processing and presentation.

Bacillus anthracis secretes two toxins into the extracellular medium during growth (30, 31). The two toxins consist of three distinct proteins that combine in a pairwise fashion (32, 33). Protective antigen (PA) combines with lethal factor (LF) or edema factor (EF). PA combined with LF constitutes lethal toxin, whereas PA combined with EF makes edema toxin (34, 35). In a simple model of intoxication, PA (83 kDa) binds to a protein receptor on the surface of cells. PA is then cleaved by a cellular protease (furin) and the amino-terminal 20-kDa fragment is released, leaving a 63-kDa fragment, PA63, bound to the cell (36–38). PA63 is able to bind either LF or EF. The toxin is then endocytosed and transported into the cell. Conversion of PA to PA63 allows formation of an oligomeric form of PA, which, after exposure to low pH in late endosomes, forms channels in cell membranes (39, 40). It is believed that these channels facilitate the delivery of LF or EF from the endosome to the cytosol of the cell (41, 42). We have previously described the internalization of fusion proteins containing anthrax toxin lethal factor and the catalytic domains of other bacterial toxins (43–45). These active fusions were used to define the amino-terminal 254 aa of LF as those necessary for uptake of the fusions (46). In this work, we have taken advantage of the efficient delivery of LF fusion proteins to the cytosol to intracellularly inoculate living cells with whole protein antigens. This system will provide the basis for new potent CTL vaccines.

MATERIALS AND METHODS

Reagents and General Procedures. Restriction endonucleases and DNA modifying enzymes were purchased from Life Technologies, Boehringer Mannheim, or New England Biolabs. Oligonucleotides were synthesized on a PCR mate (Applied Biosystems) and purified on oligonucleotide purification cartridges (Applied Biosystems). The PCR was performed with a GeneAmp kit according to the manufacturer's directions (Perkin-Elmer/Cetus). Bacterial media preparation, restriction digests, ligation, and phosphatase treatment of DNA were performed by standard protocols (47). Peptide P18IIIB was made by an automated peptide synthesizer (Applied Biosystems) and purified by HPLC liquid chromatography before use (48). Lactacystin was synthesized in the lab of E. J. Corey (Harvard University, Cambridge, MA) and was a generous gift of Jonathan Yewdell and Jack Bennink (National Institute of Allergy and Infectious Diseases, Bethesda, MD).

Plasmid Construction. The plasmid used for expressing the LF254-gp120 fusion protein in *Escherichia coli* was constructed by ligation of the pGEX-KG vector (Pharmacia) with PCR-amplified LF and gp120 gene sequences. The DNA encoding residues 1–254 of LF were amplified from plasmid pLF7 with primers that added unique *Xba*I and *Mlu*I sites on the 5' and 3' ends, respectively (49). The primers were 5'-TCT AGA TCT AGA AGC GGG CGG TCA TGG TGA TGT AGG-3' (primer 1) and 5'-GAT CTT TAA GTT CAC GCG TGG ATA GAT TTA TTT CTT G-3' (primer 2). The gene for gp120 was amplified from plasmid HXB2-env with primers that added unique restriction sites for *Mlu*I on the 5' end and *Xho*I on the 3' end of the amplified gene (43). The sequences were 5'-CCG CGT ACG CGT ATG AGA GTG AAG GAG AAA TAT CAG-3' (primer 3) and 5'-TTC GAG CTC GAG TTA TCT TTT TTC TCT CTG CAC CAC-3' (primer 4). Primer 4 introduced a stop sequence (TAA) after the gp120

coding sequence. The amplified DNA products and the pGEX-KG plasmid DNA were digested with the appropriate restriction enzymes. Vector DNA was dephosphorylated with bacterial alkaline phosphatase for 30 min. All three DNA fragments were purified from low-melting-point agarose after electrophoresis by extraction with phenol-chloroform, mixed, and ligated overnight at 16°C with T4 DNA ligase. The ligated DNA was used to transform chemically competent *E. coli* (DH5 α , high efficiency, Life Technologies). Transformed *E. coli* were selected on ampicillin containing solid media (50 μ g/ml) and screened by restriction analysis of extracted plasmid. Clones that had the expected restriction pattern were confirmed by DNA sequencing.

Expression and Purification of the LF254-gp120 Fusion Protein. The use of plasmid pGEX-KG allowed for the expression of the three-part fusion protein (GST-LF254-gp120) after induction by isopropyl-1-thio- β -D-galactopyranoside (IPTG) and rapid purification by adsorption to glutathione-Sepharose 4B. The expression and purification of GST-LF fusion proteins was described previously (44). *E. coli* strain SG12036 was transformed with the pGEX-KG-LF254-gp120 fusion vector and grown in rich media (superbroth, 100 μ g/ml of ampicillin) with shaking at 225 rpm at 37°C. When the cell density at A₆₀₀ reached 0.6–0.8, IPTG was added to a final concentration of 1 mM. After further incubation for 2 hr, the bacterial cells were pelleted by centrifugation and then resuspended in 100 mM phosphate buffer (pH 7.4), 150 mM NaCl, 1% Triton X-100, 5 mM EDTA, 5 μ g/ml leupeptin, 10 μ g/ml aprotinin, and 10 μ g/ml 4-(2-aminoethyl)-benzenesulfonylfluoride. The bacterial cells were disrupted by sonication, and the clarified extracts were applied to a glutathione-Sepharose 4B column previously equilibrated with buffer (100 mM phosphate, pH 7.4/150 mM NaCl/1% Triton X-100). The column was washed extensively, and the bound fusion protein was eluted with 10 mM glutathione in 50 mM Tris, pH 8.0/0.5 mM EDTA. The eluted protein was concentrated by ultrafiltration with a Centriprep-30 device (Amicon) and analyzed for purity by electrophoresis on nondenaturing and SDS polyacrylamide gels (Phast gels, Pharmacia). Protein concentrations were determined by the micro BCA method with BSA as a standard (Pierce).

Protective Antigen Proteins. Protective antigen was expressed in *B. anthracis* from the expression vector pYS5 and purified by established procedures (50, 51). Mutant PA molecules PA CFD and PA-D were constructed by site-directed mutagenesis and have been previously described (52).

Cell Lines. P815, a DBA/2-derived (H-2^d) mastocytoma (ATCC TIB-64) used as target cells in the CTL assay, was maintained in RPMI 1640 medium supplemented with 10% FCS. The HIV gp120-specific CTL lines 9.23.3 and α 15-12, which recognize the V3 epitope RGPGRFVTI, have been previously described (13, 48, 53). HIV gp120-specific CTL lines were derived from BALB/c spleens taken from mice previously immunized with a recombinant vaccinia virus expressing the gp160 protein and restimulated weekly. 9.23.3 CTLs were stimulated with 10 μ M P18IIIB peptide pulsed on 5×10^6 irradiated splenocytes [3,000 rads (1 rad = 0.01 Gy)] with 5×10^5 CTLs per well, in a 24-well plate containing 2 ml of a 1:1 mixture of RPMI 1640 medium and Eagle-Hanks' amino acid medium (EHAA) supplemented with L-glutamine, sodium pyruvate, nonessential amino acids, penicillin, streptomycin, 5×10^{-5} M 2-mercaptoethanol, 10% fetal calf serum, and 10% T-stim (Collaborative Biomedical Products). Stimulation of the α 15-12 CTL line used the same media mixture; however, these CTLs were restimulated with mitomycin C-treated α 15-12 cells at 5×10^5 CTLs, 5×10^5 15-12 cells, and 5×10^6 irradiated (3,000 rads) BALB/c splenocytes per well. 15-12 cells were isolated from transfected BALB/c 3T3 cells, which express the gp160 protein from the IIIB strain of HIV-1 (48).


Cytotoxic Lymphocyte Assay. ^{51}Cr release assays were performed as previously described (54). When appropriate, target cells were treated with PA (100 ng/ml) and/or LF254-gp120 fusion protein (1.0 ng/ml when used with $\alpha 15-12$ CTL or 50 ng/ml when used with 9.23.3 CTL) for 12 hr in serum-free media. After treatment, the target cells (5×10^5 cells) were labeled with 200 μCi (1 Ci = 37 GBq) of $\text{Na}_2^{51}\text{CrO}_4$ in 100 μl of RPMI 1640 for 2 hr at 37°C . In some cases, target cells were pulsed with peptide (1.0–10.0 μM) during labeling. Following labeling, targets were washed three times and added at 3,000 cells per well along with the appropriate number of effector cells in 96-well round-bottom plates. Supernatants were harvested after 6 hr and counted in an Isomedic gamma counter (ICN). The mean of triplicate samples was calculated and the percent ^{51}Cr release was determined using the following formula. Percent ^{51}Cr release = $100 \times [(\text{experimental } ^{51}\text{Cr release} - \text{control } ^{51}\text{Cr release}) / (\text{maximum } ^{51}\text{Cr release} - \text{control } ^{51}\text{Cr release})]$, where experimental ^{51}Cr release represents counts from target cells mixed with effector cells, control ^{51}Cr release represents target cells mixed with medium alone (spontaneous release), and maximum ^{51}Cr release represents counts from target cells exposed to 2.5% Triton X-100.

RESULTS

Plasmid Construction. The LF sequences required for binding to PA and translocation to the cell cytosol were genetically fused to the gp120 portion of the envelope gene from HIV-1 strain HXB2. This fusion was constructed in the pGEX-KG vector, resulting in a three-way protein fusion of the 26-kDa glutathione S-transferase domain with LF residues 1–254 and residues 1–511 of the gp160 protein. The inclusion of the Gly₅-containing linker from pGEX-KG and the alteration of a base following the *Xba*I site before the 5' end of the LF1–254 gene fragment resulted in a modified 14-residue spacer (GSPGISGGGGGILE) between the GST domain and LF1–254 (Fig. 1). The addition of the unique *Mlu*I and *Pst*I sites to the 5' end of the coding sequences for the gp120 portion of the envelope gene added four additional residues, TRILQ, between LF1–254 and gp120 (Fig. 1). The addition of residues at the amino- and carboxyl-terminal ends of LF1–254 do not appear to affect the function of the protein with respect to PA binding and protein translocation (42, 44). The resulting fusion protein has a calculated molecular mass of 114,852 Da and a calculated pI of 7.00.

Specific Cytolysis of Target Cells Sensitized with LF254-gp120 Fusion Proteins by HIV gp120-Specific CTL. The mouse mastocytoma cell line P815 was incubated with PA or PA mutants and/or LF254-gp120 fusion protein for 12 hr in serum-free conditions. The cells were then washed and labeled with ^{51}Cr for use as target cells. The labeled P815 target cells were mixed at different ratios with the effector CTL cell line 9.23.3, which recognizes the peptide epitope, RGPGRFVTI, from the V3 loop of gp120 (13, 48, 53). Killing of the target cell population was determined by measuring release of ^{51}Cr into the media (Fig. 2). Target cells treated with wild-type PA and LF254-gp120 fusion protein were efficiently recognized and lysed by the 9.23.3 CTL line. To ensure maximal lysis, effector-to-target ratios starting at 40:1 were used. Lysis was still on a plateau at an effector-to-target ratio of 10. The recognition and lysis was dependent on the presence of functional PA, demonstrating that processing of the fusion protein relies on internalization via the PA receptor. Treatment with a translocation-defective mutant PA protein (52) or without the addition of PA resulted in minimal lysis of the target cells, placing a limit on the amount of LF254-gp120 that is proteolytically degraded extracellularly and thus able to sensitize target cells without active transport into the cytosol. This implies that the LF254-gp120 fusion protein needs to gain access to the cell cytosol for processing and presentation.

A.



B.




FIG. 1. Plasmid construct and fusion protein. (A) Plasmid NA125 construct for expression of LF254-gp120 fusion protein in *E. coli*. GST, 26-kDa glutathione S-transferase domain from the parent vector pGEX-KG; LF1–254, coding sequences for residues 1–254 of mature LF; gp120, coding sequences for residues 1–511 of HXB2-env gene. (B) Expressed GST-LF254-gp120 protein. The carboxyl-terminal residue of the linker from pGEX-KG was changed from D to E, and four residues, TRILQ, were added between the LF254 and gp120 portions of the construct due to DNA manipulations. The V3 epitope, RGPGRFVTI, recognized by CTL begins at residue 807 of the fusion protein.

Treatment with PA alone, mutant PA alone, or the LF254-gp120 fusion alone did not sensitize the target cells for lysis. Target cells pulsed with P18HIB peptide served as a positive control and resulted in cell lysis when effector cells were added. To compare the relative potency of the LF254-gp120 with that of the optimal decapeptide P18-110 for sensitizing target cells for recognition by these CTL (53, 55, 56), titration experiments were performed. The P18-110 decapeptide sensitized targets for lysis in the range of 10^{-10} to 10^{-11} M, based on specific lysis $>10\%$ as a threshold for positivity, whereas the LF254-gp120 sensitized targets with as little as 4×10^{-14} M protein (data not shown). Therefore, the fusion construct was at least 2.5 logs more potent on a molar basis than the optimal binding peptide.

Lactacystin Prevents Lysis of PA/LF254-gp120-Treated Target Cells. To demonstrate the requirement for LF254-gp120 processing by the classical MHC class I pathway, we examined the ability of the specific proteasome inhibitor, lactacystin, to inhibit presentation of the V3 epitope on the cell surface. Incubation of P815 cells with 10 μM lactacystin for 45 min prior to the addition of PA and LF254-gp120 significantly decreased lysis of the target cells by the $\alpha 15-12$ CTL line (Fig. 3). Lactacystin inhibition of peptide presentation shows that the processing of the peptide epitope from the fusion protein depends on the function of the proteasome complex. This indicates that the protein fusion was delivered to the cytosol of the cell, where it interacts with the proteasome complex and precludes the role of any alternate processing pathways for presentation by the anthrax toxin LF254-gp120 fusion protein.




FIG. 2. Recognition of cells treated with gp120 anthrax toxin fusions by gp120-specific cytotoxic lymphocytes. The mouse mastocytoma cell line P815 was incubated with wild-type protective antigen (WPA) or mutant PACFD (MPA) and/or LF254-gp120 fusion proteins (50 ng/ml). P815 cells pulsed with 1.0 μ M P18111B peptide served as a control. After 12 hr, the cells were washed and the treated P815 cells were labeled with 51 Cr. Labeled P815 cells were then mixed at different ratios with the CTL cell line 9.23.3, which recognizes the peptide sequence RGPGRFVFI from gp120. Killing of the target cell population was determined by measuring release of 51 Cr into the medium (% lysis). Background lysis was determined with untreated, 51 Cr labeled, P815 cells. These data are the replicate of three experiments.

DISCUSSION

The primary goal of this study was to determine the ability of an anthrax lethal factor fusion to deliver antigenic proteins to the cell cytosol for processing and presentation with class I molecules on the cell surface. We demonstrated the ability of the LF254-gp120 fusion, in conjunction with PA, to sensitize target cells (Fig. 2). The requirement for functional PA shows that the LF254-gp120 fusion needs to gain access to the intracellular environment to be processed. Other toxins that have been used as fusions with peptide epitopes have also resulted in presentation of the epitope on the surface of antigen presenting cells in the proper context. Examples include DT, PE, pertussis toxin, and the *Bordetella pertussis* adenylate cyclase (24–27, 57). However, the processing and presentation of peptide epitopes from each of these fusion proteins do not appear to occur through the classical MHC class I processing pathway but instead via various alternate pathways.

DT was used as a delivery system with peptide epitopes from influenza virus fused to the amino-terminal end of the toxin (24). Although it was demonstrated that the peptide-toxin fusion gained access to the inside of the cell, processing and presentation of the antigens was not demonstrated. Subsequently, it was reported that DT was able to translocate additional protein domains into the cell cytosol (57, 58). Genetic fusions of an inactive DT protein with an active DT catalytic domain were translocated into the cytosol, where they were toxic. However, fusions with other protein domains had reduced abilities to deliver proteins or were unable to bind to the DT receptor and were not translocated. The entire genetic fusion appeared to be very stable in the cytosol, with no indication of any processing 6 hr after translocation, which may imply that the translocated DT or DT fusions are not accessible to the degradative machinery in the cell.

A truncated PE has also been genetically fused to influenza A peptide epitopes (25). The ability of PE-peptide antigen fusions to deliver the antigenic peptide to the MHC class I molecules on the surface of target cells was examined. Specific lysis by CTLs recognizing the influenza A epitopes within the

fusion protein was detected. However, it was determined that the PE fusions trafficked through the cell differently from native PE. In fact, blocking translocation after adding the PE fusions to cells had no effect on processing and presentation of the antigenic peptide (59). The authors proposed that a certain amount of proteolysis occurs in the endosome and that there exist, in the endosome, MHC class I molecules able to bind the processed peptide; thus, recycling of these complexes to the cell surface can mark the cell for recognition and lysis by CTLs. These PE fusions may enter a noncytosolic pathway for presenting exogenous antigens to the MHC class I molecules (60, 61).

Peptides from the lymphocytic choriomeningitis virus nucleoprotein were inserted into the S1 subunit of pertussis toxin and the pertussis adenylate cyclase (26, 27, 57, 62). These fusions specifically sensitized target cells to the appropriate CTLs. However, the dependence on the classical pathway for processing and presentation of the antigen to the MHC class I pathway was not established. The adenylate cyclase fusions were tested for their effectiveness *in vivo* and were able to generate CTLs. Production of CTLs was dependent on coinjection of the fusion protein with alum, which implies that the fusions are processed by an alternate exogenous pathway for MHC class I antigen.

In the case of LF254-gp120, we have demonstrated that delivery to the cytosol requires active PA and that presentation requires a functional proteasome complex. Use of lactacystin, a specific inhibitor for proteasomes, provides an unequivocal demonstration that these constructs enter the cytosolic degradative pathway involved in classical processing for class I MHC presentation (Fig. 3) (63). Thus, in this case a bacterial toxin fusion protein has been definitively shown to deliver a whole protein antigen to the classical cytosolic MHC class I pathway.

The anthrax lethal toxin system has several additional advantages over the other bacterial toxins for vaccine use. First, LF fusion proteins are nontoxic, because the entire catalytic domain has been deleted. Even the "detoxified" DT still retains 1/800 of the wild-type activity, which is enough to preclude its use *in vitro* or *in vivo* (24, 64). In contrast, the entire catalytic domain of LF, residues 255–776, has been deleted in LF254-gp120. Our prior studies have shown that only LF residues 1–254 are needed for efficient delivery of fused polypeptides (46). Second, LF fusions can be produced as soluble proteins in standard GST expression vectors. The binary nature of the toxin allows production of LF fusions without manipulation or alteration of the cell-binding properties of PA. LF residues 1–254 appear indifferent to addition of peptides or larger proteins at either the amino or carboxyl termini and still retain their ability to bind PA (44). These additions to LF do not appear to seriously affect the ability of PA to translocate protein into the cell cytosol. In contrast, expression and purification of the other toxin fusions require either use of special hosts capable of secretion of the modified holotoxins or refolding after purification from inclusion bodies. Third, LF fusions appear to be proficient at delivering proteins to the cell cytosol. LF-*Pseudomonas* exotoxin A fusions are 100 times more potent than wild-type PE, demonstrating that the delivery of the active domain is more efficient (43). The ability of our system to efficiently deliver antigen to the cytosol is reflected in the comparatively small amount of protein needed to achieve recognition by CTL. Both the pertussis cyclase delivery system and the pertussis toxin delivery system require 50 times more antigen on a molar basis to achieve similar recognition of target cells by CTL (26, 27). Finally, LF254 appears capable of delivering protein antigens that are large enough that hosts from varied genetic backgrounds can be expected to process the appropriate epitope from the antigen and properly present it.




FIG. 3. Inhibition of fusion protein processing by lactacystin. P815 cells were treated with 1 ng/ml of the GST-LF254-gp120 fusion protein alone and with PA (WPA) or mutant PA-D (DELPA) in the presence (B) or absence (A) of 10 μ M lactacystin. 51 Cr-labeled target cells were mixed with α 15-12 CTLs, and specific lysis was measured. Target cells were treated with the P18IIIB peptide as a positive control for cell lysis. These data are the representative of three experiments.

Preliminary *in vivo* experiments show that a single subcutaneous injection of the LF254-gp120 fusion along with PA is able to elicit a vigorous CTL response in the spleens of vaccinated animals (data not shown). We are currently refining the immunization procedure and testing the PA dependence of the *in vivo* response to confirm a cytosolic processing mechanism, as well as investigating the ability of other LF254-antigen fusions, including LF254-gp41, LF254-reverse transcriptase, and LF254-NEF, to generate an immune response.

A recent study by Ballard *et al.* (65), performed concurrently with the present study, addresses the effectiveness of *in vivo* immunization with anthrax toxin fusion constructs, using a fusion of LF with a nine-residue peptide from the listeriolysin protein of *Listeria monocytogenes*, and shows protective efficacy against infection. However, because this study used only a nonpeptide minimal fragment from the protein, which can bind directly to class I MHC molecules and does not require processing through the class I pathway, it did not address the question of whether the toxin fusion construct could introduce a whole foreign vaccine protein into the cytosolic class I MHC processing pathway and elicit appropriate degradation and presentation with a class I MHC molecule. Our present study demonstrates this critical mechanistic point and shows that whole protein antigens may be introduced in this fashion for potential use as vaccines. Whole protein antigens have the advantage that they are not limited to single epitopes that are presented by only one or a few MHC molecules and are also more susceptible to escape mutations. However, they have the disadvantage that they are usually harder to introduce into the class I MHC processing pathway without a live viral vector (17). Thus, the major advantage of anthrax toxin fusion proteins is the ability to introduce whole complex foreign proteins to this processing pathway. Thus, the Ballard *et al.* study (65) and ours complement each other in demonstrating the utility of the anthrax lethal toxin fusion system for the stimulation of a cellular immune response.

In conclusion, we have taken advantage of several unique properties of anthrax toxin to construct fusion proteins that transport large proteins into the cytosol of living cells, where they are processed through the classical pathway for loading onto MHC class I molecules and presentation to CD8 CTL. Anthrax toxin appears particularly well suited, of all the bacterial toxins, to development of safe and potent vaccines for induction of CTL immunity.

We thank Dr. E. J. Corey for synthesizing and Drs. Jonathan Yewdell and Jack Bennink for the generous gift of lactacystin.

- Robbins, J. B., Schneerson, R. & Szu, S. C. (1995) *J. Infect. Dis.* **171**, 1387-1398.
- Taylor, P. M. & Askonas, B. A. (1986) *Immunology* **58**, 417-420.
- Lin, Y. L. & Askonas, B. A. (1981) *J. Exp. Med.* **154**, 225-234.
- Lukacher, A. E., Braciale, V. L. & Braciale, T. J. (1984) *J. Exp. Med.* **160**, 814-826.
- McMichael, A. J., Gotch, F. M., Noble, G. R. & Beare, P. A. S. (1983) *N. Engl. J. Med.* **309**, 13-17.
- Byrne, J. A. & Oldstone, M. B. (1984) *J. Virol.* **51**, 682-686.
- Klavinskis, L. S., Tishon, A. & Oldstone, M. B. A. (1989) *J. Immunol.* **143**, 2013-2016.
- Klavinskis, L. S., Whitton, J. L., Joly, E. & Oldstone, M. B. A. (1990) *Virology* **178**, 393-400.
- Klavinskis, L. S., Tishon, A. & Oldstone, M. B. (1989) *J. Immunol.* **143**, 2013-2016.
- Taylor, P. M. & Askonas, B. A. (1986) *Immunology* **58**, 417-420.
- Sarin, P. S., Mora, C. A., Naylor, P. H., Markham, R., Schwartz, D., Kahn, J., Heseltine, P., Gazzard, B., Youle, M., Rios, A. & Goldstein, A. L. (1995) *Cell. Mol. Biol.* **41**, 401-407.
- van Kuyk, R., Torbett, B. E., Gulizia, R. J., Leath, S., Mosier, D. E. & Koenig, S. (1994) *J. Immunol.* **153**, 4826-4833.
- Alexander-Miller, M. A., Leggett, G. R. & Berzofsky, J. A. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 4102-4107.
- Harding, C. V. (1994) in *Cellular Proteolytic Systems*, eds. Ciechanover, A. J. & Schwartz, A. L. (Wiley, New York), pp. 163-180.
- Carbone, F. R. & Bevan, M. J. (1989) in *Fundamental Immunology*, ed. Paul, W. E. (Raven, New York), pp. 541-567.
- Townsend, A. & Bodmer, H. (1989) *Annu. Rev. Immunol.* **7**, 601-624.
- Germain, R. N. & Margulies, D. H. (1993) *Annu. Rev. Immunol.* **11**, 403-450.
- Cease, K. B. & Berzofsky, J. A. (1994) *Annu. Rev. Immunol.* **12**, 923-989.
- Takahashi, H., Takeshita, T., Morein, B., Putney, S. D., Germain, R. N. & Berzofsky, J. A. (1990) *Nature (London)* **344**, 873-875.
- Ulmer, J. B., Donnelly, J. J., Parker, S. E., Rhodes, G. H., Felgner, P. L., Dworki, V. J., Gromkowski, S. H., Deck, R. R., DeWitt, C. M., Friedman, A., Hawe, L. A., Leander, K. R., Martinez, D., Perry, H. C., Shiver, J. W., Montgomery, D. L. & Liu, M. A. (1993) *Science* **259**, 1745-1749.
- White, W. I., Cassatt, D. R., Madsen, J., Burke, S. J., Woods, R. M., Wassef, N. M., Alving, C. R. & Koenig, S. (1995) *Vaccine* **13**, 1111-1122.
- Daniel, M. D., Kirchhoff, F., Czajak, S. C., Sehgal, P. K. & Desrosiers, R. C. (1992) *Science* **258**, 1938-1941.

23. Baba, T. W., Trichel, A. M., An, L., Liska, V., Martin, L. N., Murphey-Corb, M. & Ruprecht, R. M. (1996) *Science* **272**, 1486-1489.
24. Stenmark, H., Moskaug, J. O., Madshus, I. H., Sandvig, K. & Olsnes, S. (1991) *J. Cell Biol.* **113**, 1025-1032.
25. Donnelly, J. J., Ulmer, J. B., Hawe, L. A., Friedman, A., Shi, X. P., Leander, K. R., Shiver, J. W., Oliff, A. I., Martinez, D. & Montgomery, D. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 3530-3534.
26. Carbonetti, N., Tishon, A., Oldstone, M. & Rappuoli, R. (1995) *Abstr. Annu. Meet. Am. Soc. Microbiol.* **95**, 295.
27. Sebo, P., Fayolle, C., d'Andria, O., Ladant, D., Leclerc, C. & Ullmann, A. (1995) *Infect. Immunol.* **63**, 3851-3857.
28. Del Val, M., Schlicht, H.-J., Ruppert, T., Reddehase, M. J. & Koszinowski, U. H. (1991) *Cell* **66**, 1145-1153.
29. Hahn, Y. S., Braciale, V. & Braciale, T. J. (1991) *J. Exp. Med.* **174**, 733-736.
30. Smith, H. & Keppie, J. (1954) *Nature (London)* **173**, 869-870.
31. Smith, H. & Stanley, J. L. (1962) *J. Gen. Microbiol.* **29**, 517-521.
32. Smith, H. & Stoner, H. B. (1967) *Fed. Proc.* **26**, 1554-1557.
33. Leppla, S. (1995) in *Bacterial Toxins and Virulence Factors in Disease: Handbook of Natural Toxins, Vol 8*, eds. Moss, J., Iglewski, B., Vaughan, M. & Tu, A. (Dekker, New York), pp. 543-572.
34. Friedlander, A. M. (1986) *J. Biol. Chem.* **261**, 7123-7126.
35. Leppla, S. H. (1982) *Proc. Natl. Acad. Sci. USA* **79**, 3162-3166.
36. Leppla, S. H., Klimpel, K. R. & Arora, N. (1994) in *Molecular Mechanisms of Bacterial Virulence*, eds. Kado, C. I. & Crosa, J. H. (Kluwer, Dordrecht, The Netherlands), pp. 127-139.
37. Klimpel, K. R., Molloy, S. S., Thomas, G. & Leppla, S. H. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 10277-10281.
38. Novak, J. M., Stein, M. P., Little, S. F., Leppla, S. H. & Friedlander, A. M. (1992) *J. Biol. Chem.* **267**, 17186-17193.
39. Blaustein, R. O., Koehler, T. M., Collier, R. J. & Finkelstein, A. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 2209-2213.
40. Milne, J. C. & Collier, R. J. (1993) *Mol. Microbiol.* **10**, 647-653.
41. Milne, J. C., Furlong, D., Hanna, P. C., Wall, J. S. & Collier, R. J. (1994) *J. Biol. Chem.* **269**, 20607-20612.
42. Milne, J. C., Blanke, S. R., Hanna, P. C. & Collier, R. J. (1995) *Mol. Microbiol.* **15**, 661-666.
43. Arora, N., Klimpel, K. R., Singh, Y. & Leppla, S. H. (1992) *J. Biol. Chem.* **267**, 15542-15548.
44. Arora, N. & Leppla, S. H. (1994) *Infect. Immunol.* **62**, 4955-4961.
45. Arora, N., Williamson, L. C., Leppla, S. H. & Halpern, J. L. (1994) *J. Biol. Chem.* **269**, 26165-26171.
46. Arora, N. & Leppla, S. H. (1993) *J. Biol. Chem.* **268**, 3334-3341.
47. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Cold Spring Harbor, NY).
48. Takahashi, H., Cohen, J., Hosmalin, A., Cease, K. B., Houghten, R., Cornette, J., DeLisi, C., Moss, B., Germain, R. N. & Berzofsky, J. A. (1988) *Proc. Natl. Acad. Sci. USA* **85**, 3105-3109.
49. Robertson, D. L. & Leppla, S. H. (1986) *Gene* **44**, 71-78.
50. Singh, Y., Chaudhary, V. K. & Leppla, S. H. (1989) *J. Biol. Chem.* **264**, 19103-19107.
51. Leppla, S. H. (1988) in *Methods in Enzymology, Vol. 165*, ed. Harshman, S. (Academic, Orlando, FL), pp. 103-116.
52. Singh, Y., Klimpel, K. R., Arora, N., Sharma, M. & Leppla, S. H. (1994) *J. Biol. Chem.* **269**, 29039-29046.
53. Shirai, M., Pendleton, C. D. & Berzofsky, J. A. (1992) *J. Immunol.* **148**, 1657-1667.
54. Alexander, M. A., Damico, C. A., Wietes, K. M., Hansen, T. H. & Connolly, J. M. (1991) *J. Exp. Med.* **173**, 849-858.
55. Kozlowski, S., Corr, M., Takeshita, T., Boyd, L. F., Pendleton, C. D., Germain, R. N., Berzofsky, J. A. & Margulies, D. H. (1992) *J. Exp. Med.* **175**, 1417-1422.
56. Takeshita, T., Takahashi, H., Kozlowski, S., Ahlers, J. D., Pendleton, C. D., Moore, R. L., Nakagawa, Y., Yokomuro, K., Fox, B. S., Margulies, D. H. & Berzofsky, J. A. (1995) *J. Immunol.* **154**, 1973-1986.
57. Madshus, I. H., Olsnes, S. & Stenmark, H. (1992) *Infect. Immunol.* **60**, 3296-3302.
58. Madshus, I. H., Stenmark, H., Sandvig, K. & Olsnes, S. (1991) *J. Biol. Chem.* **266**, 17446-17453.
59. Ulmer, J. B., Donnelly, J. J. & Liu, M. A. (1994) *Eur. J. Immunol.* **24**, 1590-1596.
60. Kovacsics-Bankowski, M. & Rock, K. L. (1995) *Science* **267**, 243-246.
61. Rock, K. L. (1996) *Immunol. Today* **17**, 131-137.
62. Fayolle, C., Sebo, P., Ladant, D., Ullmann, A. & Leclerc, C. (1996) *J. Immunol.* **156**, 4697-4706.
63. Fenteany, G., Standaert, R. F., Lane, W. S., Choi, S., Corey, E. J. & Schreiber, S. L. (1995) *Science* **268**, 726-731.
64. Barbieri, J. T. & Collier, R. J. (1987) *Infect. Immunol.* **55**, 1647-1651.
65. Ballard, J. D., Collier, R. J. & Starnbach, M. N. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 12531-12534.



UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office
Address: COMMISSIONER OF PATENTS AND TRADEMARKS
Washington, D.C. 20231
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
08/937,276	09/15/1997	KURT KLIMPEL	15280-290100	1627

7590

03/08/2002

KENNETH A WEBER
TOWNSEND AND TOWNSEND AND CREW
TWO EMBARCADERO CENTER 8TH FLOOR
SAN FRANCISCO, CA 941113834

EXAMINER

SCHWADRON, RONALD B

ART UNIT PAPER NUMBER

1644

DATE MAILED: 03/08/2002

Response Due 6-8-02 MNR

Please find below and/or attached an Office communication concerning this application or proceeding.

Office Action Summary

Application No.

08/937,276

Applicant(s)

Klimpel et al.

Examiner

Ron Schwadron

Art Unit

1644

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

Period for Reply

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136 (a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If the period for reply specified above is less than thirty (30) days, a reply within the statutory minimum of thirty (30) days will be considered timely.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133).
- Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

Status

- 1) ☒ Responsive to communication(s) filed on 8/30/2001 and 12/17/2001.
- 2a) ☐ This action is **FINAL**. 2b) ☒ This action is non-final.
- 3) ☐ Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11; 453 O.G. 213.

Disposition of Claims

- 4) ☒ Claim(s) 9-19 is/are pending in the application.
- 4a) Of the above, claim(s) _____ is/are withdrawn from consideration.
- 5) ☐ Claim(s) _____ is/are allowed.
- 6) ☒ Claim(s) 9-19 is/are rejected.
- 7) ☐ Claim(s) _____ is/are objected to.
- 8) ☐ Claims _____ are subject to restriction and/or election requirement.

Application Papers

- 9) ☐ The specification is objected to by the Examiner.
- 10) ☐ The drawing(s) filed on _____ is/are objected to by the Examiner.
- 11) ☐ The proposed drawing correction filed on _____ is: a) ☐ approved b) ☐ disapproved.
- 12) ☐ The oath or declaration is objected to by the Examiner.

Priority under 35 U.S.C. § 119

- 13) ☐ Acknowledgement is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d).
- a) ☐ All b) ☐ Some* c) ☐ None of:
- ☐ Certified copies of the priority documents have been received.
 - ☐ Certified copies of the priority documents have been received in Application No. _____.
 - ☐ Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).

*See the attached detailed Office action for a list of the certified copies not received.

- 14) ☐ Acknowledgement is made of a claim for domestic priority under 35 U.S.C. § 119(e).

Attachment(s)

- 15) ☐ Notice of References Cited (PTO-892) 18) ☐ Interview Summary (PTO-413) Paper No(s). _____
- 16) ☐ Notice of Draftsperson's Patent Drawing Review (PTO-948) 19) ☐ Notice of Informal Patent Application (PTO-152)
- 17) ☐ Information Disclosure Statement(s) (PTO-1449) Paper No(s). _____ 20) ☐ Other:

1. Claims 9-19 are under consideration. Claims 1-7,20-28 have been canceled.

RESPONSE TO APPLICANTS ARGUMENTS

2. The following is a quotation of the first paragraph of 35 U.S.C. 112:

The specification shall contain a written description of the invention, and of the manner and process of making and using it, in such full, clear, concise, and exact terms as to enable any person skilled in the art to which it pertains, or with which it is most nearly connected, to make and use the same and shall set forth the best mode contemplated by the inventor of carrying out his invention.

3. Claims 9-19 are rejected under 35 U.S.C. 112, first paragraph, because the specification, while being enabling for the claimed invention which uses an APABP construct which contains amino acids 1-254 of LF is not enabling for the claimed invention which uses intact LF. Arora et al. (1993) discloses that intact LF causes cell death when administered to cells (see page 3334, first column). Thus, the claimed invention which uses intact LF could not be used to induce an immune response to CTL epitopes because it would kill the treated cells. The specification does not enable any person skilled in the art to which it pertains, or with which it is most nearly connected, to practice the invention commensurate in scope with these claims.

4. Claims 9-19 are rejected under 35 U.S.C. § 112, second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject matter which applicant regards as the invention.

Claim 9 is indefinite in the recitation of "anthrax protective antigen binding protein". The specification page 7, lines 1 and 2, defines said term as "contains the PA binding site of LF". The specification page 7, lines 21-23 defines LF as a 90 kDa protein derived from *B. anthracis*. Said protein is known in the art. The indefinite issue arises in that page 7, lines 23-24 recites "The description of LF includes binary toxin functional equivalents such as protein Ia of *C. perfringens*.". It is unclear from this sentence whether LF is limited to the 90 kDa protein derived from *B. anthracis* or whether LF encompasses includes binary

toxin functional equivalents because it is unclear whether "The description of LF includes" refers to the definition of LF or has some other meaning. If the term is taken as encompassing "binary toxin functional equivalents", then it is unclear what this term means or encompasses because it is not defined in the specification and it has no art recognized meaning. It is unclear as to what properties or set of properties would be encompassed by a "binary toxin functional equivalent".

5. Claims 9-19 are rejected under 35 U.S.C. 112, first paragraph, as containing subject matter which was not described in the specification in such a way as to reasonably convey to one skilled in the relevant art that the inventor(s), at the time the application was filed, had possession of the claimed invention.

The specification does not provide adequate written description of the claimed invention. The legal standard for sufficiency of a patent's (or a specification's) written description is whether that description "reasonably conveys to the artisan that the inventor had possession at that time of the . . . claimed subject matter", *Vas-Cath, Inc. V. Mahurkar*, 19 U.S.P.Q.2d 1111 (Fed. Cir. 1991). In the instant case, the specification does not convey to the artisan that the applicant had possession at the time of invention of the claimed inventions.

As per paragraph 4 of the instant Office Action, instant claims encompass use of a "binary toxin functional equivalents". With the exception of the particular LF derived fragment derived from the first 254 amino acids of the 90 kDA protein of *B. anthracis* disclosed in the specification, the skilled artisan cannot envision the detailed structure of the encompassed "binary toxin functional equivalents" and therefore conception is not achieved until reduction to practice has occurred, regardless of the complexity or simplicity of the method of isolation. For example, it is clear that the peptide recited in the claims encompasses a vast number of potential analogs with varying amino acid sequences, yet there is no disclosure in the specification of what amino acid sequence would be used other than the specific example disclosed in the specification. Adequate written description requires more than a mere statement that it is part of the invention and a reference to a potential method of isolating it. In the instant application, the amino acid itself or isolated protein is required. See *Fiers v. Revel*, 25 USPQ 2d 1601 at 1606 (CAFC 1993) and *Amgen Inc. V. Chugai Pharmaceutical Co. Ltd.*, 18 USPQ2d 1016.

In view of the aforementioned problems regarding description of the claimed invention, the specification does not provide an adequate written description of the invention claimed herein. See *The Regents of the University of California v. Eli Lilly and Company*, 43 USPQ2d 1398, 1404-7 (Fed. Cir. 1997). In *University of California v. Eli Lilly and Co.*, 39 U.S.P.Q.2d 1225 (Fed. Cir. 1995) the inventors claimed a genus of DNA species encoding insulin in different vertebrates or mammals, but had only described a single species of cDNA which encoded rat insulin. The court held that only the nucleic acids species described in the specification (i.e. nucleic acids encoding rat insulin) met the description requirement and that the inventors were not entitled to a claim encompassing a genus of nucleic acids encoding insulin from other vertebrates, mammals or humans, *id.* at 1240. The Federal Circuit has held that if an inventor is "unable to envision the detailed constitution of a gene so as to distinguish it from other materials. . .conception has not been achieved until reduction to practice has occurred", *Amgen, Inc. v. Chugai Pharmaceutical Co, Ltd.*, 18 U.S.P.Q.2d 1016 (Fed. Cir. 1991). Attention is also directed to the decision of *The Regents of the University of California v. Eli Lilly and Company* (CAFC, July 1997) wherein is stated:

"The description requirement of the patent statute requires a description of an invention, not an indication of a result that one might achieve if one made that invention. See *In re Wilder*, 736 F.2d 1516, 222 USPQ 369, 372-373 (Fed. Cir. 1984) (affirming rejection because the specification does "little more than outlin[e] goals appellants hope the claimed invention achieves and the problems the invention will hopefully ameliorate."). Accordingly, naming a type of material generally known to exist, in the absence of knowledge as to what that material consists of, is not a description of that material. Thus, as we have previously held, a cDNA is not defined or described by the mere name "cDNA," even if accompanied by the name of the protein that it encodes, but requires a kind of specificity usually achieved by means of the recitation of the sequence of nucleotides that make up the cDNA." See *Fiers*, 984 F.2d at 1171, 25 USPQ2d at 1606.

7. The rejection of claims 9-19 under 35 U.S.C. 103(a) as being unpatentable over *Milne et al.* (*Mol. Microbiology* 15:651, 1995), *Arora et al.* (*J. Biol. Chemistry* 268:3334, 1993) or *Leppla et al.* US Patent 5,591,631 in view of EP 0 532 090A2 (issued March

Art Unit: 1644

3, 1991) and Donnelly et al . (PNAS 90:3530:1993) is withdrawn in view of applicants arguments and the Berzofsky declaration filed 8/30/2001.

8. No claim is allowed.

9. Papers related to this application may be submitted to Group 1600 by facsimile transmission. The faxing of such papers must conform with the notice published in the Official Gazette, 1096 OG 30 (November 15, 1989). Papers should be faxed to Group 1600 at (703) 308-4242.

10. Any inquiry concerning this communication or earlier communications from the examiner should be directed to Dr. Ron Schwadron whose telephone number is (703) 308-4680. The examiner can normally be reached Monday through Thursday from 7:30 to 6:00. A message may be left on the examiners voice mail service. If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, Ms. Christina Chan can be reached on (703) 308-3973. Any inquiry of a general nature or relating to the status of this application should be directed to the Group 1600 receptionist whose telephone number is (703) 308-0196.



RONALD B. SCHWADRON
PRIMARY EXAMINER
GROUP 1800 *1600*

Ron Schwadron, Ph.D.

Primary Examiner

Art Unit 1644

Attachment for PTO-948 (Rev. 03/01, or earlier)

6/18/01

The below text replaces the pre-printed text under the heading, "Information on How to Effect Drawing Changes," on the back of the PTO-948 (Rev. 03/01, or earlier) form.

INFORMATION ON HOW TO EFFECT DRAWING CHANGES

1. Correction of Informalities -- 37 CFR 1.85

New corrected drawings must be filed with the changes incorporated therein. Identifying indicia, if provided, should include the title of the invention, inventor's name, and application number, or docket number (if any) if an application number has not been assigned to the application. If this information is provided, it must be placed on the front of each sheet and centered within the top margin. If corrected drawings are required in a Notice of Allowability (PTOL-37), the new drawings **MUST** be filed within the **THREE MONTH** shortened statutory period set for reply in the Notice of Allowability. Extensions of time may **NOT** be obtained under the provisions of 37 CFR 1.136(a) or (b) for filing the corrected drawings after the mailing of a Notice of Allowability. The drawings should be filed as a separate paper with a transmittal letter addressed to the Official Draftsperson.

2. Corrections other than Informalities Noted by Draftsperson on form PTO-948.

All changes to the drawings, other than informalities noted by the Draftsperson, **MUST** be made in the same manner as above except that, normally, a highlighted (preferably red ink) sketch of the changes to be incorporated into the new drawings **MUST** be approved by the examiner before the application will be allowed. No changes will be permitted to be made, other than correction of informalities, unless the examiner has approved the proposed changes.

Timing of Corrections

Applicant is required to submit the drawing corrections within the time period set in the attached Office communication. See 37 CFR 1.85(a).

Failure to take corrective action within the set period will result in **ABANDONMENT** of the application.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKewed/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.